



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 305/14	A1	(11) International Publication Number: WO 99/09021 (43) International Publication Date: 25 February 1999 (25.02.99)								
<p>(21) International Application Number: PCT/US98/17016</p> <p>(22) International Filing Date: 17 August 1998 (17.08.98)</p> <p>(30) Priority Data:</p> <table border="0"> <tr> <td>60/056,000</td> <td>18 August 1997 (18.08.97)</td> <td>US</td> </tr> <tr> <td>60/081,265</td> <td>9 April 1998 (09.04.98)</td> <td>US</td> </tr> <tr> <td>09/063,477</td> <td>20 April 1998 (20.04.98)</td> <td>US</td> </tr> </table> <p>(71) Applicant: FLORIDA STATE UNIVERSITY [US/US]; 100 Sliger Building, Innovation Park, Tallahassee, FL 32306-2763 (US).</p> <p>(72) Inventors: HOLTON, Robert, A.; 7125 Upland Glade, Tallahassee, FL 32312 (US). ZHANG, Zhuming; 36 Hawthorne Place #4S, Montclair, NJ 07042-3246 (US). CLARK, Paul, A.; 1701 W. Pensacola Street #201, Tallahassee, FL 32304 (US).</p> <p>(74) Agents: HEILEK, Edward, J. et al.; Senniger, Powers, Leavitt & Roedel, 16th floor, One Metropolitan Square, St. Louis, MO 63102 (US).</p>	60/056,000	18 August 1997 (18.08.97)	US	60/081,265	9 April 1998 (09.04.98)	US	09/063,477	20 April 1998 (20.04.98)	US	<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
60/056,000	18 August 1997 (18.08.97)	US								
60/081,265	9 April 1998 (09.04.98)	US								
09/063,477	20 April 1998 (20.04.98)	US								
<p>(54) Title: PROCESS FOR SELECTIVE DERIVATIZATION OF TAXANES</p> <p>(57) Abstract</p> <p>Processes for the preparation of taxol and other taxanes through selective derivatization of the C(7) and C(10) hydroxyl groups of 10-DAB, particularly a novel process using a new strategy in which the C(10) hydroxyl group is protected or derivatized prior to the C(7) hydroxyl group; and the provision of C(7) and C(10) derivatized 10-DAB compounds.</p>										

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.


AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

PROCESS FOR SELECTIVE DERIVATIZATION OF TAXANES

BACKGROUND OF THE INVENTION

The present invention is directed, in general, to a process for the preparation of taxol and other taxanes, and in particular, to such a process in which the C(7) or C(10) hydroxyl group of a taxane is selectively derivatized.

10-DAB (1), which is extracted from the needles of *taxus baccata* L., the English yew, has become a key starting material in the production of taxol and Taxotere, both of which are potent anticancer agents. Conversion of 10-DAB to taxol, Taxotere® and other taxanes having antitumor activity requires protection or derivatization of the C(7) and C(10) hydroxyl groups followed by esterification of the C(13) hydroxyl group to attach an appropriate side chain at that position.



- 1 R₁₀ = R₇ = H
2 R₁₀ = H, R₇ = TES

Until now, strategies for the preparation of taxol and taxol analogs were based upon the observation of Senilh et al. (C.R. Acad. Sci. Paris, II, 1981, 293, 501) that the relative reactivity of the four hydroxyl groups of 10-DAB toward acetic anhydride in pyridine is
5 C(7)-OH > C(10)-OH > C(13)-OH > C(1)-OH. Denis, et. al. reported (J. Am. Chem. Soc., 1988, 110, 5917) selective silylation of the C(7) hydroxyl group of 10-DAB with triethylsilyl chloride in pyridine to give
10 7-triethylsilyl-10-deacetyl baccatin (III) (2) in 85% yield. Based upon these reports, in those processes in which differentiation of the C(7) and C(10) hydroxyl groups is required (e.g., preparation of taxol from 10-DAB), the C(7) hydroxyl group must be protected (or
15 derivatized) before the C(10) hydroxyl group is protected or derivatized. For example, taxol may be prepared by treating 10-DAB with triethylsilyl chloride to protect the C(7) hydroxyl group, acetylating the C(10) hydroxyl group, attaching the side chain by esterification of the
20 C(13) hydroxyl group, and, finally, removal of protecting groups.

It is known that taxanes having various substituents bonded to either the C(10) or the C(7) oxygens show anticancer activity. To provide for more efficient
25 synthesis of these materials, it would be useful to have methods which permit more efficient and more highly selective protection or derivatization of the C(10) and the C(7) hydroxyl groups.

SUMMARY OF THE INVENTION

30 Among the objects of the present invention, therefore, is the provision of highly efficient processes for the preparation of taxol and other taxanes through selective derivatization of the C(7) group or the C(10) hydroxyl group of 10-DAB and other taxanes, particularly


a process in which the C(10) hydroxyl group is protected or derivatized prior to the C(7) hydroxyl group; and the provision of C(7) or C(10) derivatized taxanes.

Briefly, therefore, the present invention is
 5 directed to a process for the acylation of the C(10) hydroxy group of a taxane. The process comprises forming a reaction mixture containing the taxane and an acylating agent which contains less than one equivalent of a base for each equivalent of taxane, and allowing the taxane to
 10 react with the acylating agent to form a C(10) acylated taxane.

The present invention is further directed to a process for the silylation of the C(10) hydroxy group of a taxane having a C(10) hydroxy group. The process
 15 comprises treating the taxane with a silylamide or a bissilylamide to form a C(10) silylated taxane.

The present invention is further directed to a process for converting the C(7) hydroxy group of a 10-acyloxy-7-hydroxytaxane to an acetal or ketal. The
 20 process comprises treating the 10-acyloxy-7-hydroxytaxane with a ketalizing agent in the presence of an acid catalyst to form a C(10) ketalized taxane.

The present invention is further directed to a taxane having the structure:



25

wherein

M is a metal or comprises ammonium;

R₁ is hydrogen, hydroxy, protected hydroxy, or together with R₁₄ or R₂ forms a carbonate;

R_2 is keto, $-OT_2$, acyloxy, or together with R_1 forms a carbonate;

R_4 is $-OT_4$ or acyloxy;

R_7 is $-OSiR_JR_KR_L$;

5 R_9 is hydrogen, keto, $-OT_9$, or acyloxy;

R_{10} is hydrogen, keto, $-OT_{10}$, or acyloxy;

R_{13} is hydroxy, protected hydroxy, keto, or $MO-$;

R_{14} is hydrogen, $-OT_{14}$, acyloxy, or together with R_1 forms a carbonate;

10 R_J , R_K , R_L are independently hydrocarbyl, substituted hydrocarbyl, or heteroaryl, provided, however, if each of R_J , R_K and R_L are alkyl, at least one of R_J , R_K and R_L comprises a carbon skeleton having at least four carbon atoms; and

15 T_2 , T_4 , T_9 , T_{10} , and T_{14} are independently hydrogen or hydroxy protecting group.

Other objects and features of this invention will be in part apparent and in part pointed out hereinafter.


DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

20 Among other things, the present invention enables the selective derivatization of the C(10) hydroxyl group of a taxane without first protecting the C(7) hydroxyl group. Stated another way, it has been discovered that the reactivities previously reported for the C(7) and
25 C(10) hydroxyl groups can be reversed, that is, the reactivity of the C(10) hydroxyl group becomes greater than the reactivity of the C(7) hydroxyl group under certain conditions.

30 Although the present invention may be used to selectively derivatize a taxane having a hydroxy group at C(7) or C(10), it offers particular advantages in the selective derivatization of taxanes having hydroxy groups

5

at C(7) and C(10), i.e., 7,10-dihydroxy taxanes. In general, 7,10-dihydroxytaxanes which may be selectively derivatized in accordance with the present invention correspond to the following structure:



3

wherein


R₁ is hydrogen, hydroxy, protected hydroxy, or together with R₁₄ or R₂ forms a carbonate;

10 R₂ is keto, -OT₂, acyloxy, or together with R₁ forms a carbonate;

R₄ is -OT₄ or acyloxy;

R₉ is hydrogen, keto, -OT₉, or acyloxy;

R₁₃ is hydroxy, protected hydroxy, keto, or



15

R₁₄ is hydrogen, -OT₁₄, acyloxy or together with R₁ forms a carbonate;

T₂, T₄, T₉, and T₁₄ are independently hydrogen or hydroxy protecting group;

20 X₁ is -OX₆, -SX₇, or -NX₈X₉;

X₂ is hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl;

X₃ and X₄ are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl;

25 X₅ is -X₁₀, -OX₁₀, -SX₁₀, -NX₈X₁₀, or -SO₂X₁₁;

X_6 is hydrocarbyl, substituted hydrocarbyl, heteroaryl, hydroxy protecting group or a functional group which increases the water solubility of the taxane derivative;

5 X_7 is hydrocarbyl, substituted hydrocarbyl, heteroaryl, or sulfhydryl protecting group;

X_8 is hydrogen, hydrocarbyl, or substituted hydrocarbyl;

X_9 is an amino protecting group;

10 X_{10} is hydrocarbyl, substituted hydrocarbyl, or heteroaryl;

X_{11} is hydrocarbyl, substituted hydrocarbyl, heteroaryl, $-OX_{10}$, or $-NX_8X_{14}$; and



X_{14} is hydrogen, hydrocarbyl, substituted
15 hydrocarbyl, or heteroaryl.

Selective C(10) Derivatization

In accordance with the process of the present invention, it has been discovered that the C(10) hydroxyl group of a taxane can be selectively acylated in the
20 absence of a base, preferably in the absence of an amine base. Preferably, therefore, amine bases such as pyridine, triethylamine, dimethylaminopyridine and 2,6-lutidine, if present at all, are present in the reaction mixture in relatively low concentration. Stated another
25 way, if a base is present in the reaction mixture, the molar ratio of the amine base to the taxane is preferably less than 1:1, more preferably less than 10:1, and most preferably less than 100:1.

Acylating agents which may be used for the selective
30 acylation of the C(10) hydroxyl group of a taxane include anhydrides, dicarbonates, thiodicarbonates, and isocyanates. In general, the anhydrides, dicarbonates, and thiodicarbonates correspond to structure 4 and the isocyanates correspond to structure 5:

7



wherein R^1 is $-\text{OR}^a$, $-\text{SR}^a$, or R^a ; R^2 is $-\text{OC}(\text{O})\text{R}^b$, $-\text{OC}(\text{O})\text{OR}^b$, $-\text{OC}(\text{O})\text{SR}^b$, $-\text{OPOR}^b\text{R}^c$, or $-\text{OS}(\text{O})_2\text{R}^b$; R^3 is hydrocarbyl, substituted hydrocarbyl, or heteroaryl; and R^a , R^b , R^c are independently hydrocarbyl, substituted hydrocarbyl, or heteroaryl. For example, suitable carboxylic acid anhydride acylating agents include acetic anhydride, chloroacetic anhydride, propionic anhydride, benzoic anhydride, and other carboxylic acid anhydrides containing substituted or unsubstituted hydrocarbyl or heteroaryl moieties; suitable dicarbonate acylating reagents include dibenzyl dicarbonate, diallyl dicarbonate, dipropyl dicarbonate, and other dicarbonates containing substituted or unsubstituted hydrocarbyl or heteroaryl moieties; and suitable isocyanate acylating agents include phenyl isocyanate, and other isocyanates containing substituted or unsubstituted hydrocarbyl or heteroaryl moieties. In addition, although the anhydrides, dicarbonates, and thiodicarbonates used as acylating agents may be mixed, it is generally preferred that they be symmetrical; that is, R^1 and R^2 are selected such that the molecule is symmetrical (e.g., if R^1 is R^a , R^2 is $-\text{OC}(\text{O})\text{R}^b$ with R^a being the same as R^b).




While the acylation of the C(10) hydroxy group of the taxane will proceed at an adequate rate for many acylating agents, it has been discovered that the reaction rate may be increased by including a Lewis acid in the reaction mixture. The concentration of the Lewis acid appears not to be narrowly critical; experimental evidence obtained to date suggests it may be present in either a stoichiometric or a catalytic amount. In general, Lewis acids which may be used include triflates and halides of elements of groups IB, IIB, IIIB, IVB, VB,

VIB, VIIB, VIII, IIIA, IVA, lanthanides, and actinides of the Periodic Table (American Chemical Society format). Preferred Lewis acids include zinc chloride, stannic chloride, cerium trichloride, cuprous chloride, lanthanum trichloride, dysprosium trichloride, and ytterbium trichloride. Zinc chloride or cerium trichloride is particularly preferred when the acylating agent is an anhydride or dicarbonate. Cuprous chloride is particularly preferred when the acylating agent is an isocyanate.



The solvent for the selective acylation is preferably an ethereal solvent such as tetrahydrofuran. Alternatively, however, other solvents such as ether or dimethoxyethane may be used.

The temperature at which the C(10) selective acylation is carried out is not narrowly critical. In general, however, it is preferably carried out at room temperature or higher in order for the reaction to proceed at a sufficiently high rate.


For purposes of illustration, acylating reactions involving dibenzyl dicarbonate, diallyl dicarbonate, acetic anhydride, chloroacetic anhydride and phenyl isocyanate are illustrated in Reaction Schemes 1 through 5 below. In this series of reaction schemes, the taxane which is selectively acylated at the C(10) position is 10-deacetylbaccatin III. It should be understood, however, that these reaction schemes are merely illustrative and that other taxanes having a C(10) hydroxy group, in general, and other 7,10-dihydroxytaxanes, in particular, may be selectively acylated with these and other acylating agents in accordance with the present invention.

Scheme 1Scheme 25 Scheme 3

10

Scheme 4Scheme 5

- 5 In another aspect of the present invention, the C(10) hydroxyl group of a taxane may be selectively silylated. In general, the silylating agent is selected from the group consisting of silylamides and bissilylamides. Preferred silylamides and bissilyamides
- 10 correspond to structures 6 and 7, respectively:



- wherein R_D , R_E , R_F , R_G , and R_H are independently hydrocarbyl, substituted hydrocarbyl, or heteroaryl.
- 15 Preferably, the silylating agents are selected from the group consisting of tri(hydrocarbyl)silyl-trifluoromethylacetamides and bis tri(hydrocarbyl)-

silyltrifluoromethylacetamides, with the hydrocarbonyl moiety being substituted or unsubstituted alkyl or aryl. For example, the preferred silylamides and bissilylamides include N,O-bis-(trimethylsilyl) trifluoroacetamide, N,O-bis-(triethylsilyl)trifluoroacetamide, N-methyl-N-triethylsilyltrifluoroacetamide, and N,O-bis(t-butylldimethylsilyl)trifluoroacetamide.



The silylating agents may be used either alone or in combination with a catalytic amount of a base such as an alkali metal base. Alkali metal amides, such as lithium amide catalysts, in general, and lithium hexamethyldisilazide, in particular, are preferred.

The solvent for the selective silylation reaction is preferably an ethereal solvent such as tetrahydrofuran. Alternatively, however, other solvents such as ether or dimethoxyethane may be used.

The temperature at which the C(10) selective silylation is carried out is not narrowly critical. In general, however, it is carried out at 0 °C or greater.

Selective C(10) silylation reactions involving N,O-bis(trimethylsilyl)trifluoroacetamide and N,O-bis(triethylsilyl)trifluoroacetamide are illustrated in Reaction Schemes 6 and 7 below. In these reaction schemes, the taxane which is selectively silylated at the C(10) position is 10-deacetylbaaccatin III. It should be understood, however, that these reaction schemes are merely illustrative and that other taxanes, having a C(10) hydroxy group, in general, and other 7,10-dihydroxytaxanes, in particular, may be selectively silylated with these and other silylating agents in accordance with the present invention.

12

Scheme 6Scheme 7

- 5 After the C(10) hydroxyl group of a
 7,10-dihydroxytaxane has been derivatized as described
 herein, the C(7) hydroxyl group can readily be protected
 or otherwise derivatized selectively in the presence of
 the C(1) and C(13) hydroxyl groups (and a C(14) hydroxy
 10 group, if present).

Selective C(7) Derivatization


- 15 Selective acylation of the C(7) hydroxyl group of a
 C(10) acylated or silylated taxane can be achieved using
 any of a variety of common acylating agents including,
 but not limited to, substituted and unsubstituted
 carboxylic acid derivatives, e.g., carboxylic acid
 halides, anhydrides, dicarbonates, isocyanates and
 haloformates. For example, the C(7) hydroxyl group of
 baccatin III, 10-acyl-10-deacetylbaccatin III or

10-triethylsilyl-10-deacetyl baccatin III can be selectively acylated with dibenzyl dicarbonate, diallyl dicarbonate, 2,2,2-trichloroethyl chloroformate, benzyl chloroformate or another common acylating agent.




5 In general, acylation of the C(7) hydroxy group of a C(10) acylated or silylated taxane are more efficient and more selective than are C(7) acylations of a 7,10-dihydroxy taxane such as 10-DAB, i.e., once the C(10) hydroxyl group has been acylated or silylated, there is a
10 significant difference in the reactivity of the remaining C(7), C(13), and C(1) hydroxyl groups (and the C(14) hydroxyl group, if present). These acylation reactions may optionally be carried out in the presence or absence of a base.

15 Examples of selective C(7) acylation of a taxane having an acylated or silylated C(10) hydroxy group are shown in Reaction Schemes 8 through 11. In these reaction schemes, the taxane which is selectively acylated at the C(7) position is baccatin III or
20 10-triethylsilyl-10-deacetyl baccatin III. It should be understood, however, that these reaction schemes are merely illustrative and that taxanes having other acyl and silyl moieties at C(10) as well as other substituents at other taxane ring positions may be selectively
25 acylated at C(7) with these and other acylating agents in accordance with the present invention.

Scheme 8



14

Scheme 9Scheme 105 Scheme 11




Alternatively, the C(7) hydroxyl group of a C(10) acylated taxane derivative can be selectively protected using any of a variety of hydroxy protecting groups, such as acetal, ketal, silyl, and removable acyl protecting groups. For example, the C(7) hydroxyl group may be silylated using any of a variety of common silylating agents including, but not limited to,

tri(hydrocarbyl)silyl halides and tri(hydrocarbyl)silyl triflates. The hydrocarbyl moieties of these compounds may be substituted or unsubstituted and preferably are substituted or unsubstituted alkyl or aryl. For example, the C(7) hydroxyl group of baccatin III can be selectively silylated using silylating agents such as tribenzylsilyl chloride, trimethylsilyl chloride, triethylsilyl chloride, dimethyl isopropylsilyl chloride, dimethyl phenylsilyl chloride, and the like.


In general, silylations of the C(7) hydroxy group of a C(10) acylated taxanes are more efficient and more selective than are silylations of a 7,10-dihydroxy taxane such as 10-DAB, i.e., once the C(10) hydroxyl group has been acylated, there is a significant difference in the reactivity of the remaining C(7), C(13), and C(1) hydroxyl groups (and the C(14) hydroxyl group, if present). The C(7) silylation reaction may be carried out under a wide range of conditions, including in the presence or absence of an amine base.

Examples of selective C(7) silylation of C(10) acylated taxanes are shown in Reaction Schemes 12 through 15. In these reaction schemes, the taxane which is selectively silylated at the C(7) position is baccatin III or another C(10)-acyloxy derivative of 10-deacetyl baccatin III. It should be understood, however, that these reaction schemes are merely illustrative and that other taxanes may be selectively silylated with these and other silylating agents in accordance with the present invention.

16

Scheme 12Scheme 135 Scheme 14

Scheme 15




Alternatively, the C(7) hydroxyl group of C(10) acylated taxanes can be selectively protected using any of a variety of common reagents including, but not limited to, simple acetals, ketals and vinyl ethers, in the presence of an acid catalyst. These reagents (whether acetal, ketal, vinyl ether or otherwise) are referred to herein as "ketalizing agents" and are described in "Protective Groups in Organic Synthesis" by T. W. Greene, John Wiley and Sons, 1981. The acid catalyst used may be an organic or inorganic acid, such as toluenesulfonic acid or camphorsulfonic acid, in at least a catalytic amount. For example, the C(7) hydroxyl group of baccatin III can be selectively ketalized using 2-methoxy propene. Other suitable reagents for the preparation of acetals and ketals include methyl vinyl ether, ethyl vinyl ether, tetrahydropyran, and the like.

Selective ketalization of the C(7) substituent of a C(10) acylated taxane is more efficient and more selective than it is with 10-DAB, i.e., once the C(10) hydroxyl group has been acylated, there is a large difference in the reactivity of the remaining C(7), C(13), and C(1) hydroxyl groups (and the C(14) hydroxyl group, if present).

An example of selective formation of a C(7) ketal from baccatin III is illustrated in Reaction Scheme 16. It should be understood, however, that this reaction

scheme is merely illustrative and that other taxanes may be selectively ketalized with this and other ketalizing agents in accordance with the present invention.



Scheme 16





Under appropriate conditions, the C(7) hydroxyl group of a taxane further comprising a C(10) hydroxyl group can be selectively silylated. Advantageously, these silylations are not limited to silyl groups bearing alkyl substituents having three carbons or less.

In general, the C(7) hydroxyl group of a taxane can be selectively silylated with a silylating agent which includes the $-\text{SiR}_j\text{R}_k\text{R}_l$ moiety wherein R_j , R_k and R_l are independently substituted or unsubstituted hydrocarbyl or heteroaryl, provided that any substituents are other than hydroxyl. In one embodiment of the present invention, if each of R_j , R_k and R_l is alkyl, then at least one of R_j , R_k , and R_l comprises a carbon skeleton (i.e., carbon chain or ring(s)) having at least four carbon atoms. Suitable silylating agents include silyl halides and silyl triflates, for example, tri(hydrocarbyl)silyl halides and tri(hydrocarbyl)silyl triflates. The hydrocarbyl substituents of these silylating agents may be substituted or unsubstituted and preferably are substituted or unsubstituted alkyl or aryl.

The selective silylation of the C(7) hydroxy group may be carried out in a solvent, such as dimethyl formamide ("DMF") or pyridine and in the presence of an amine base, such as imidazole or pyridine. Reaction Schemes 17 - 20 illustrate the silylation of the C(7) hydroxy group of 10-DAB in high yield by treating 10-DAB with t-butyldimethylsilyl chloride, tribenzylsilyl chloride, dimethyl-isopropylsilyl chloride, and dimethylphenylsilyl chloride, respectively. Silylation under these conditions was surprising in view of the report by Denis, et. al. (*J. Am. Chem. Soc.*, 1988, 110, 5917) that selective formation of 7-TBS-10-DAB was not possible.



Scheme 17Scheme 18

Scheme 19Scheme 20

5 The process of the present invention can also be
 used to protect the C(7) and C(10) hydroxy groups of a
 7,10-dihydroxytaxane with different silyl protecting
 groups. By selecting groups which can be removed under
 different conditions, the C(7) and C(10) hydroxy groups
 10 can be separately accessed for derivatization. These
 reactions, therefore, increase the flexibility of the
 overall process and, enable a higher yield for many of
 the individual protecting reactions relative to the yield
 obtained using currently available processes. For
 15 example, the triethylsilyl protecting group is more
 readily removed from C(10) than is the
 t-butyldimethylsilyl protecting group from C(7) and the
 dimethylphenylsilyl protecting group is more readily
 removed from C(7) than is the t-butyldimethylsilyl


21

protecting group from C(10). The preparation of 7-t-butyldimethylsilyl-10-triethylsilyl-10-DAB and 7-dimethylphenylsilyl-10-t-butyldimethylsilyl-10-DAB are illustrated in Reaction Schemes 21 and 22.

5 Scheme 21Scheme 22

The methods disclosed herein may be used in connection with a large number of different taxanes obtained from natural or synthetic sources to prepare a wide variety of taxane intermediates which may then be further derivatized. For example, the methods of the present invention may be effectively used to protect the C(7) and/or C(10) hydroxy functional group prior to the coupling reaction between a C(13) side chain precursor and a taxane to introduce a C(13) β -amido ester side chain, and also prior to the reactions for preparing taxanes having alternative substituents at various locations on the taxane nucleus.

The attachment of a C(13) side chain precursor to a taxane may be carried out by various known techniques. For example, a side chain precursor such as an appropriately substituted β -lactam, oxazoline, oxazolidine carboxylic acid, oxazolidine carboxylic acid anhydride, or isoserine derivative may be reacted with a tricyclic or tetracyclic taxane having a C(13) hydroxy, metallic oxide or ammonium oxide substituent to form compounds having a β -amido ester substituent at C(13) as described, for example, Taxol: Science and Applications, M. Suffness, editor, CRC Press (Boca Rotan, FL) 1995, Chapter V, pages 97-121. For example, the synthesis of taxol from 10-DAB is illustrated in reaction scheme 23. It should be noted that while a β -lactam and 10-DAB are used in this reaction scheme, other side chain precursors and other taxanes could be substituted therefor without departing from the present invention.


Scheme 23

The process illustrated in Reaction Scheme 23 is significantly more efficient than any other currently known process, due to the high yields and selectivity of the cerium trichloride catalyzed acetylation of the C(10) hydroxyl group of 10-DAB and the subsequent silylation of the C(7) hydroxyl group. The synthesis proceeds in four steps and 89% overall yield.

Reaction schemes 24 and 25 illustrate the preparation of taxanes having substituents appended to the C(7) hydroxyl group and a free C(10) hydroxyl group. The method of the current invention provides flexibility so that the substituent attached to the C(7) hydroxyl group can be put in place either before or after attachment of the C(13) side chain.

Reaction scheme 24 outlines the preparation of a taxane which has been found to be a potent chemotherapeutic radiosensitizer, illustrating attachment of the substituent at the C(7) hydroxyl group before introduction of the C(13) side chain. According to the process of reaction scheme 7, 10-DAB is first converted to 10-TES-10-DAB. The C(7) hydroxyl group is then converted to an intermediate imidazolidine by treatment with carbonyl diimidazole, and the intermediate imidazolidine subsequently reacts, without isolation, with metronidazole alcohol to provide 7-metro-10-TES-10-DAB. Coupling of 7-metro-10-TES-10-DAB with a β -lactam to introduce the side chain at C(13) is followed by removal of the TES groups at C(10) and C(2') by treatment with HF and pyridine.

Scheme 24




25

Reaction scheme 25 outlines the preparation of a taxane useful in identifying proteins which form bioconjugates with taxanes. It illustrates a protocol for attachment of a substituent at the C(7) hydroxyl group after introduction of the C(13) side chain.

5 According to the processes of reaction schemes 7 and 11, 10-DAB is first converted to 7-*p*-nitrobenzyloxycarbonyl-10-TES-10-DAB. The C(13) side chain is attached employing a TES protected β -lactam, and the

10 *p*-nitrobenzyloxycarbonyl protecting group is then selectively removed by treatment with hydrogen and a palladium catalyst, producing 2',10-(bis)-TES-taxotere. The C(7) hydroxyl group then reacts with carbonyl diimidazole and the derived imidazolide is treated with

15 1,4-diamino butane to give a primary amine. Reaction of the primary amine with the hydroxysuccinimide ester of biotin completes the attachment of the biotinamide group at C(7). Finally, treatment with HF in pyridine solution removes the TES protecting groups at C(10) and C(2').

Scheme 25

The protected taxane derivatives or the intermediates or starting materials used in the preparation of such protected taxane derivatives can be further modified to provide for alternative substituents at various positions of the taxane.

Taxanes having C(2) and/or C(4) substituents other than benzyloxy and acetoxy, respectively, can be prepared from baccatin III, 10-DAB and other taxanes as more fully described in PCT Patent Application WO 94/01223. In general, the C(2) and C(4) acyloxy substituents are treated with lithium aluminum hydride or another suitable reducing agent to form hydroxy groups at C(2) and C(4) which may then be reacted, for example, with carboxylic acid halides (optionally after protection of the C(2) hydroxy group together with the C(1) hydroxy group with a 1,2-carbonate protecting group) to obtain the desired C(2) and C(4) derivatives.

Taxanes having C(7) substituents other than hydroxy and acyloxy as described herein can be prepared from baccatin III, 10-DAB, and other taxanes as more fully described in PCT Patent Application WO 94/17050. For example, a C(7) xanthate may be subjected to tin hydride reduction to yield the corresponding C(7) dihydro taxane. Alternatively, C(7) fluoro-substituted taxanes can be prepared by treatment of C(13)-triethylsilyl-protected baccatin III with 2-chloro-1,1,2-trifluoroethylamine at room temperature in THF solution. Other baccatin derivatives with a free C(7) hydroxyl group behave similarly. Alternatively, 7-chloro baccatin III can be prepared by treatment of baccatin III with methanesulfonyl chloride and triethylamine in methylene chloride solution containing an excess of triethylamine hydrochloride.

Taxanes having C(9) substituents other than keto can be prepared from baccatin III, 10-DAB and other taxanes as more fully described in PCT Patent Application WO


94/20088. In general, the C(9) keto substituent of the taxane is selectively reduced to yield the corresponding C(9) β -hydroxy derivative with a borohydride, preferably tetrabutylammonium borohydride (Bu_4NBH_4) or triacetoxy-borohydride. The C(9) β -hydroxy derivative can then be protected at C(7) with a hydroxy protecting group and the C(9) hydroxy group can be acylated following the methods described herein for acylation of the C(7) hydroxy group. Alternatively, reaction of 7-protected-9 β -hydroxy derivative with KH causes the acetate group (or other acyloxy group) to migrate from C(10) to C(9) and the hydroxy group to migrate from C(9) to C(10), thereby yielding a 10-desacetyl derivative, which can be acylated as described elsewhere herein.

Taxanes having C(10) substituents other than hydroxy, acyloxy or protected hydroxy as described herein may be prepared as more fully described in PCT Patent Application WO 94/15599 and other literature references. For example, taxanes having a C(10) keto substituent can be prepared by oxidation of 10-desacetyl taxanes. Taxanes which are dihydro substituted at C(10) can be prepared by reacting a C(10) hydroxy or acyloxy substituted taxane with samarium diiodide.

Taxanes having a C(14) substituent other than hydrogen may also be prepared. The starting material for these compounds may be, for example, a hydroxylated taxane (14-hydroxy-10-deacetylbaaccatin III) which has been discovered in an extract of yew needles (C&EN, p 36-37, April 12, 1993). Derivatives of this hydroxylated taxane having the various C(2), C(4), C(7), C(9), C(10), C3' and C5' functional groups described above may also be prepared by using this hydroxylated taxane. In addition, the C(14) hydroxy group together with the C(1) hydroxy group of 10-DAB can be converted to a 1,2-carbonate as


described in C&EN or it may be converted to a variety of esters or other functional groups as otherwise described herein in connection with the C(2), C(4), C(9) and C(10) substituents.

- 5 The process of the present invention thus enables the preparation of taxanes having the following structure:



wherein

- 10 M comprises ammonium or is a metal;
 R₁ is hydrogen, hydroxy, protected hydroxy, or
 together with R₁₄ or R₂ forms a carbonate;
 R₂ is keto, -OT₂, acyloxy, or together with R₁ forms
 a carbonate;
 15 R₄ is -OT₄ or acyloxy;
 R₇ is hydrogen, halogen, -OT₇, or acyloxy;
 R₉ is hydrogen, keto, -OT₉, or acyloxy;
 R₁₀ is hydrogen, keto, -OT₁₀, or acyloxy;
 R₇, R₉, and R₁₀ independently have the alpha or beta
 20 stereochemical configuration;
 R₁₃ is hydroxy, protected hydroxy, keto, MO- or



 R₁₄ is hydrogen, -OT₁₄, acyloxy, or together with R₁
 forms a carbonate;

T_2 , T_4 , T_7 , T_9 , T_{10} and T_{14} are independently hydrogen or hydroxy protecting group;

X_1 is $-OX_6$, $-SX_7$, or $-NX_8X_9$;

X_2 is hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl;

X_3 and X_4 are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl;

X_5 is $-X_{10}$, $-OX_{10}$, $-SX_{10}$, $-NX_8X_{10}$, or $-SO_2X_{11}$;

X_6 is hydrogen, hydrocarbyl, substituted hydrocarbyl, heteroaryl, hydroxy protecting group, or a functional group which increases the water solubility of the taxane derivative;

X_7 is hydrocarbyl, substituted hydrocarbyl, heteroaryl, or sulfhydryl protecting group;

X_8 is hydrogen, hydrocarbyl, or substituted hydrocarbyl;

X_9 is an amino protecting group;

X_{10} is hydrocarbyl, substituted hydrocarbyl, or heteroaryl;


X_{11} is hydrocarbyl, substituted hydrocarbyl, heteroaryl, $-OX_{10}$, or $-NX_8X_{14}$;

X_{14} is hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl.

In one embodiment of the present invention, the substituents of the taxane (other than the C(7), C(10) and C(13) substituents) correspond to the substituents present on baccatin III or 10-DAB. That is, R_{14} is hydrogen, R_9 is keto, R_4 is acetoxy, R_2 is benzoyloxy, and R_1 is hydroxy. In other embodiments, the taxane has a structure which differs from that of taxol or Taxotere[®] with respect to the C(13) side chain and at least one other substituent. For example, R_{14} may be hydroxy; R_2 may be hydroxy, $-OCOZ_2$ or $-OCOOZ_{22}$ wherein Z_2 is hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl and Z_{22} is hydrocarbyl, substituted hydrocarbyl, or heteroaryl; R_4 may be hydroxy, $-OCOZ_4$ or $-OCOOZ_{44}$ wherein


Z_4 is hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl and Z_{44} is hydrocarbyl, substituted hydrocarbyl, or heteroaryl; R_7 may be hydrogen, hydroxy, $-\text{OCOZ}_7$, or $-\text{OCOOZ}_{77}$, wherein Z_7 is hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl and Z_{77} is hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl, R_9 may be hydrogen, hydroxy, $-\text{OCOZ}_9$, or $-\text{OCOOZ}_{99}$, wherein Z_9 is hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl and Z_{99} is hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl, and R_{10} may be hydrogen, hydroxy, $-\text{OCOZ}_{10}$ or $-\text{OCOOZ}_{1010}$ wherein Z_{10} is hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl and Z_{1010} is hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl.

In a preferred embodiment, the taxane has the formula



wherein P_{10} is acyl, said acyl comprising at least three carbon atoms or two carbon atoms and a nitrogen, oxygen or sulfur atom. Stated another way, $-\text{OP}_{10}$ is other than acetoxy. More preferably, P_{10} is $-(\text{C}=\text{O})\text{R}_A$, $-(\text{C}=\text{O})\text{OR}_B$, or $-(\text{C}=\text{O})\text{NR}_C$ wherein R_A is substituted or unsubstituted hydrocarbyl or heteroaryl, said unsubstituted hydrocarbyl comprising at least two carbon atoms; R_B and R_C are independently substituted or unsubstituted hydrocarbyl. Still more preferably, R_A is substituted or unsubstituted alkyl or aryl, said unsubstituted alkyl comprising at least two carbon atoms; and R_B and R_C are independently substituted or unsubstituted alkyl or aryl.

In another embodiment of the invention, the taxane has the formula



wherein P_7 and P_{10} are independently substituted or
 5 unsubstituted acyl. In this embodiment, P_7 and P_{10} are preferably different.

Definitions

As used herein, the terms "selective" and "selective
 10 derivatization" shall mean that the desired product is preferentially formed over any other by-product. Preferably, the desired product is present in a molar ratio of at least 9:1 relative to any other by-product and, more preferably, is present in a molar ratio of at least 20:1 relative to any other by-product.

15 In addition, "Ph" means phenyl; "Bz" means benzoyl; "Bn" means benzyl; "Me" means methyl; "Et" means ethyl; "iPr" means isopropyl; "tBu" and "t-Bu" means tert-butyl; "Ac" means acetyl; "TES" means triethylsilyl; "TMS" means trimethylsilyl; "TBS" means $\text{Me}_2\text{t-BuSi-}$; "CDI" means
 20 carbonyl diimidazole; "BOM" means benzyloxymethyl; "DBU" means diazabicycloundecane; "DMAP" means p-dimethylamino pyridine; "LHMDS" or "LiHMDS" means lithium hexamethyldisilazide; "DMF" means dimethylformamide; "10-DAB" means 10-desacetylbaccatin III; "Cbz" means
 25 benzyloxycarbonyl; "Alloc" means allyloxycarbonyl; "THF" means tetrahydrofuran; "BOC" means benzyloxycarbonyl; "PNB" means para-nitrobenzyl; "Troc" means 2,2,2-trichloroethoxycarbonyl; "EtOAc" means ethyl acetate;

"THF" means tetrahydrofuran; "protected hydroxyl" means -OP wherein P is a hydroxyl protecting group; and "hydroxyl protecting group" includes, but is not limited to, acetals having two to ten carbons, ketals having two to ten carbons, and ethers, such as methyl, t-butyl, benzyl, p-methoxybenzyl, p-nitrobenzyl, allyl, trityl, methoxymethyl, methoxyethoxymethyl, ethoxyethyl, methoxypropyl, tetrahydropyranyl, tetrahydrothiopyranyl; and trialkylsilyl ethers such as trimethylsilyl ether, triethylsilyl ether, dimethylarylsilyl ether, triisopropylsilyl ether and t-butyl dimethylsilyl ether; esters such as benzoyl, acetyl, phenylacetyl, formyl, mono-, di-, and trihaloacetyl such as chloroacetyl, dichloroacetyl, trichloroacetyl, trifluoroacetyl; and carbonates including but not limited to alkyl carbonates having from one to six carbon atoms such as methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl; isobutyl, and n-pentyl; alkyl carbonates having from one to six carbon atoms and substituted with one or more halogen atoms such as 2,2,2-trichloroethoxymethyl and 2,2,2-trichloroethyl; alkenyl carbonates having from two to six carbon atoms such as vinyl and allyl; cycloalkyl carbonates having from three to six carbon atoms such as cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl; and phenyl or benzyl carbonates optionally substituted on the ring with one or more C₁₋₆ alkoxy, or nitro. Other hydroxyl protecting groups may be found in "Protective Groups in Organic Synthesis" by T. W. Greene, John Wiley and Sons, 1981, and Second Edition, 1991.

The "hydrocarbon" and "hydrocarbyl" moieties described herein are organic compounds or radicals consisting exclusively of the elements carbon and hydrogen. These moieties include alkyl, alkenyl, alkynyl, and aryl moieties. These moieties also include alkyl, alkenyl, alkynyl, and aryl moieties substituted

with other aliphatic or cyclic hydrocarbyl groups, and include alkaryl, alkenaryl and alkynaryl. Preferably, these moieties comprise 1 to 20 carbon atoms.

The alkyl groups described herein are preferably
5 lower alkyl containing from one to six carbon atoms in the principal chain and up to 20 carbon atoms. They may be straight, branched chain or cyclic and include methyl, ethyl, propyl, isopropyl, butyl, hexyl and the like. They may be substituted with aliphatic or cyclic
10 hydrocarbyl radicals.

The alkenyl groups described herein are preferably lower alkenyl containing from two to six carbon atoms in the principal chain and up to 20 carbon atoms. They may be straight or branched chain and include ethenyl,
15 propenyl, isopropenyl, butenyl, isobutenyl, hexenyl, and the like. They may be substituted with aliphatic or cyclic hydrocarbyl radicals.

The alkynyl groups described herein are preferably lower alkynyl containing from two to six carbon atoms in the principal chain and up to 20 carbon atoms. They may be straight or branched chain and include ethynyl, propynyl, butynyl, isobutynyl, hexynyl, and the like. They may be substituted with aliphatic or cyclic
20 hydrocarbyl radicals.

25 The aryl moieties described herein contain from 6 to 20 carbon atoms and include phenyl. They may be hydrocarbyl substituted with the various substituents defined herein. Phenyl is the more preferred aryl.


The heteroaryl moieties described are heterocyclic
30 compounds or radicals which are analogous to aromatic compounds or radicals and which contain a total of 5 to 20 atoms, usually 5 or 6 ring atoms, and at least one atom other than carbon, such as furyl, thienyl, pyridyl and the like. The heteroaryl moieties may be substituted
35 with hydrocarbyl, heterosubstituted hydrocarbyl or hetero-atom containing substituents with the hetero-atoms

being selected from the group consisting of nitrogen, oxygen, silicon, phosphorous, boron, sulfur, and halogens. These substituents include hydroxy; lower alkoxy such as methoxy, ethoxy, butoxy; halogen such as
 5 chloro or fluoro; ethers; acetals; ketals; esters; heteroaryl such as furyl or thienyl; alkanoxy; acyl; acyloxy; nitro; amino; and amido.

The substituted hydrocarbyl moieties described herein are hydrocarbyl moieties which are substituted
 10 with at least one atom other than carbon and hydrogen, including moieties in which a carbon chain atom is substituted with a hetero atom such as nitrogen, oxygen, silicon, phosphorous, boron, sulfur, or a halogen atom. These substituents include hydroxy; lower alkoxy such as
 15 methoxy, ethoxy, butoxy; halogen such as chloro or fluoro; ethers; acetals; ketals; esters; heteroaryl such as furyl or thienyl; alkanoxy; acyl; acyloxy; nitro; amino; and amido.

The acyl moieties and the acyloxy moieties described
 20 herein contain hydrocarbyl, substituted hydrocarbyl or heteroaryl moieties. In general, they have the formulas $-C(O)G$ and $-OC(O)G$, respectively, wherein G is substituted or unsubstituted hydrocarbyl, hydrocarbyloxy, hydrocarbylamino, hydrocarbylthio or heteroaryl.

25 The ketal moieties described herein have the general formula




wherein X^{31} , X^{32} , X^{33} and X^{34} are independently hydrocarbyl, substituted hydrocarbyl or heteroaryl moieties. They may be optionally substituted with the various substituents
 35 defined herein. The ketal moieties are preferably substituted or unsubstituted alkyl or alkenyl, and more preferably substituted or unsubstituted lower (C_1-C_6)

alkyl. These ketal moieties additionally may encompass sugars or substituted sugars and include ketal moieties prepared from sugars or substituted sugars such as glucose and xylose. When a ketal moiety is incorporated into a taxane of the present invention as a C(7) hydroxy protecting group, then either X³¹ or X³² represents the taxane moiety.


The acetal moieties described herein have the general formula

10



wherein X³¹, X³² and X³³ are independently hydrocarbyl, substituted hydrocarbyl or heteroaryl moieties. They may be optionally substituted with the various substituents defined herein other than hydroxyl. The acetal moieties are preferably substituted or unsubstituted alkyl or alkenyl, and more preferably substituted or unsubstituted lower (C₁-C₆) alkyl. These acetal moieties additionally may encompass sugars or substituted sugars and include acetal moieties prepared from sugars or substituted sugars such as glucose and xylose. When an acetal moiety is incorporated into a taxane of the present invention as a C(7) hydroxy protecting group, then either X³¹ or X³² represents the taxane moiety.

The term "taxane" as used herein, denotes compounds containing the A, B and C rings (with numbering of the ring positions shown herein):



The following examples illustrate the invention.

Example 1

A. Selective acylation of a C(10) hydroxyl group:

10-Cbz-10-DAB. To a solution of 10-DAB (30 mg,
5 0.055 mmol) in THF (1 mL) at room temperature was added
dibenzyl pyrocarbonate (320 mg, 1.1 mmol, 20 equiv) under
N₂. The reaction mixture was stirred at room temperature
for 24 h. EtOAc (10 mL) was added, and the solution was
quickly filtered through a short column of silica gel.
10 The silica gel was washed with EtOAc (100 mL), and the
solution was concentrated under reduced pressure. The
residue was purified by flash column chromatography using
EtOAc: hexanes (1:1) as the eluent and dried in vacuo
overnight to give 10-cbz-10-DAB as a colorless solid:
15 yield, 37 mg (98%). mp 205-206 °C; $[\alpha]_D^{25}$ -63° (CHCl₃, c =
0.41); ¹H NMR (400 MHz, CDCl₃) δ 1.11(s, 3 H, Me17),
1.13(s, 3 H, Me16), 1.58(s, 1 H, 1-OH), 1.71(s, 3 H,
Me19), 1.89(ddd, J = 14.7, 10.9, 2.3 Hz, 1 H, H6b),
2.00(d, J = 5.1 Hz, 1 H, 13-OH), 2.08(d, J = 1.0, 3 H,
20 Me18), 2.28(s, 3 H, 4-Ac), 2.30(m, 2 H, H14a, H14b),
2.43(d, J = 4.1 Hz, 1 H, 7-OH), 2.58(ddd, J = 14.7, 9.6,
6.6 Hz, 1 H, H6a), 3.88(d, J = 6.9 Hz, 1 H, H3), 4.19(d,
J = 8.6 Hz, 1 H, H20b), 4.31(d, J = 8.6 Hz, 1 H, H20a),
4.44(ddd, J = 10.9, 6.6, 4.1 Hz, 1 H, H7), 4.89(m, 1 H,
25 H13), 4.98(dd, J = 9.6, 2.3 Hz, 1 H, H5), 5.23(d, J =
12.1, 1 H, CHH'OC(O)), 5.26(d, J = 12.1, 1 H, CHH'OC(O)),
5.65(d, J = 6.9 Hz, 1 H, H2), 6.19(s, 1 H, H10),
7.35-7.44(m, 5 H, PhCH₂O), 7.48(dd, J = 8.1, 7.6 Hz, 2 H,
benzoate, m), 7.60(tt, J = 7.6, 1.0 Hz, 1 H, benzoate,
30 p), 8.11(d, J = 8.1, 1.0 Hz, 2 H, benzoate, o) ppm. ¹³C
NMR (75 MHz, CDCl₃) δ 9.1(Me(19)), 15.3(Me(18)),
20.7(4-Ac), 22.3, 26.7(Me16, Me17), 35.5(C(6)),
38.6(C(14)), 42.5(C(15)), 46.1(C(3)), 58.7(C(8)),
67.9(C(13)), 70.5(OCH₂Ph), 72.2, 75.0, 76.5(C(7), C(2),
35 C(20)), 79.0, 79.1 (C(1), C(10)), 80.9(C(4)), 84.5(C(5)),

128.6, 128.8, 129.7, 130.3, 131.9, 133.8(OCH₂Ph, benzoate), 135.1(C(11)), 147.5(C(12)), 155.6(OC(O)O), 167.4(benzoate), 171.0(4-Ac), 204.7(C(9))ppm. Anal. Calcd for C₃₇H₄₂O₁₂. 1/2H₂O: C, 64.62; H, 6.30. Found: C, 64.34; H, 6.31.

10-Alloc-10-DAB. To a solution of 10-DAB (30 mg, 0.055 mmol) in THF (1 mL) at room temperature was added diallyl pyrocarbonate (366 mL, 2.2 mmol, 40 equiv) under N₂. The reaction mixture was stirred at room temperature for 48 h. TLC analysis indicated the presence of the desired product along with unreacted starting material. EtOAc (20 mL) was added, and the solution was quickly filtered through a short column of silica gel. The silica gel was washed with EtOAc (100 mL), and the solution was concentrated. The residue was purified by flash column chromatography using EtOAc: hexanes (1:1) as the eluent and dried in vacuo overnight to give 10-alloc-10-DAB as a colorless solid: yield, 23 mg (67%, 95% at the conversion of 70%). The recovered 10-DAB, 9 mg (30%). 10-alloc-10-DAB: mp 201-203 °C; [α]_D²⁰ -81° (CHCl₃, c = 0.53); ¹H NMR (400 MHz, CDCl₃) δ 1.11(s, 3 H, Me17), 1.12(s, 3 H, Me16), 1.60(s, 1 H, 1-OH), 1.69(s, 3 H, Me19), 1.87(ddd, J = 14.7, 11.0, 2.1 Hz, 1 H, H6b), 2.05(d, J = 5.1 Hz, 1 H, 13-OH), 2.08(d, J = 1.2, 3 H, Me18), 2.28(s, 3 H, 4-Ac), 2.29(m, 2 H, H14a, H14b), 2.47(d, J = 4.2 Hz, 1 H, 7-OH), 2.57(ddd, J = 14.7, 9.6, 6.7 Hz, 1 H, H6a), 3.86(d, J = 7.0 Hz, 1 H, H3), 4.16(d, J = 8.4 Hz, 1 H, H20b), 4.31(d, J = 8.4 Hz, 1 H, H20a), 4.44(ddd, J = 11.0, 6.7, 4.2 Hz, 1 H, H7), 4.70(br d, J = 5.9 Hz, 2 H, CHH'=CHCH₂O), 4.90(m, 1 H, H13), 4.97(dd, J = 9.6, 2.1 Hz, 1 H, H5), 5.32(dd, J = 10.4, 1.2 Hz, 1 H, CHH'=CHCH₂O), 5.42(dd, J = 17.2, 1.2 Hz, 1 H, CHH'=CHCH₂O), 5.63(d, J = 7.0 Hz, 1 H, H2), 5.98(ddt, J = 17.2, 10.4, 5.9 Hz, 1 H, CHH'=CHCH₂O), 6.16(s, 1 H, H10), 7.48(dd, J = 8.1, 7.5 Hz, 2 H, benzoate, m), 7.60(tt, J =

7.5, 1.2 Hz, 1 H, benzoate, p), 8.11(d, J = 8.1, 1.2 Hz, 2 H, benzoate, o) ppm; ^{13}C NMR (75 MHz, CDCl_3) δ 9.1(Me(19)), 15.3(Me(18)), 20.7(4-Ac), 22.3, 26.7(Me16, Me17), 35.5(C(6)), 38.6(C(14)), 42.5(C(15)), 46.1(C(3)), 58.7(C(8)), 67.9(C(13)), 69.3($\text{CH}_2=\text{CHCH}_2\text{O}$), 72.1, 75.0, 76.5(C(7), C(2), C(20)), 79.0, 79.1(C(1), C(10)), 80.9(C(4)), 84.5(C(5)), 119.6($\text{CH}_2=\text{CHCH}_2\text{O}$), 128.8, 129.7, 130.3, 133.8(benzoate), 131.4, 131.9($\text{CH}_2=\text{CHCH}_2\text{O}$, C(11)), 147.5(C(12)), 155.4(OC(O)O), 167.4(benzoate), 170.9(4-Ac), 204.7(C(9)) ppm. Anal. Calcd for $\text{C}_{33}\text{H}_{40}\text{O}_{12}$: C, 63.05; H, 6.41. Found: C, 62.77; H, 6.48.

B. Selective acylation of a C(10) hydroxyl group using ZnCl_2 :

baccatin III. To a solution of 10-DAB (100 mg, 0.184 mmol) in THF (6 mL) at room temperature was added a mixture of acetic anhydride (6.5 mL) and ZnCl_2 / THF solution (0.5 M, 726 mL, 0.368 mmol, 2 equiv) under N_2 . The reaction mixture was stirred at room temperature for 4 h. Then the reaction mixture was diluted with EtOAc (100 mL), washed with saturated aqueous NaHCO_3 solution (40 mLx3), brine. The organic layer was dried over Na_2SO_4 , concentrated under reduced pressure. The residue was purified by flash column chromatography using EtOAc: hexanes (1:1) as the eluent and dried in vacuo to give *baccatin III* as a colorless solid: yield, 100 mg (93%). mp 237-238 °C dec (ref 236-238 °C dec); $[\alpha]_{\text{D}}^{25} -63^\circ$ (CH_3OH , c = 0.45) (ref $[\alpha]_{\text{D}}^{25} -54^\circ$, CH_3OH); ^1H NMR (400 MHz, CDCl_3) δ 1.11(s, 6 H, Me16, Me17), 1.61(s, 1 H, 1-OH), 1.67(s, 3 H, Me19), 1.87(ddd, J = 14.7, 10.9, 2.1 Hz, 1 H, H6b), 2.05(d, J = 3.8 Hz, 1 H, 13-OH), 2.05(s, 3 H, Me18), 2.24(s, 3 H, 10-Ac), 2.28(s, 3 H, 4-Ac), 2.30(m, 2 H, H14a, H14b), 2.47(d, J = 4.2 Hz, 1 H, 7-OH), 2.57(ddd, J = 14.7, 9.4, 6.7 Hz, 1 H, H6a), 3.89(d, J = 7.0 Hz, 1 H, H3), 4.16(d, J = 8.4 Hz, 1 H, H20b), 4.31(d, J = 8.4 Hz, 1 H, H20a), 4.47(ddd, J = 10.9, 6.7, 4.2 Hz, 1 H, H7),

40

4.90(m, 1 H, H13), 4.99(dd, J = 9.4, 2.1 Hz, 1 H, H5),
5.63(d, J = 7.0 Hz, 1 H, H2), 6.33(s, 1 H, H10), 7.48(dd,
J = 7.8, 7.8 Hz, 2 H, benzoate, m), 7.61(dd, J = 7.8, 7.4
Hz, 1 H, benzoate, p), 8.11(d, J = 7.4 Hz, 2 H, benzoate,
5 o)ppm. ¹³C NMR (100 MHz, CDCl₃) δ 9.4(Me(19)),
15.6(Me(18)), 20.9(4-Ac, 10-Ac), 22.6, 27.0(Me16, Me17),
35.6(C(6)), 38.6(C(14)), 42.7(C(15)), 46.1(C(3)),
58.8(C(8)), 68.0(C(13)), 72.3, 75.0, 76.2, 76.4(C(7),
C(2), C(10), C(20)), 79.1(C(1)), 80.9(C(4)), 84.5(C(5)),
10 128.6, 129.4, 130.1, 133.7(benzoate), 132.0(C(11)),
146.3(C(12)), 167.1(benzoate), 170.7, 171.3(10-Ac, 4-Ac),
204.1(C(9))ppm.

10-Chloroacetyl-10-DAB. To a solution of 10-DAB
(116 mg, 0.21 mmol) in THF (3 mL) at room temperature was
15 added a mixture of chloroacetic anhydride (2.8 g, 16.3
mmol, 78 equiv) and ZnCl₂/ THF solution (0.5 M, 0.85 mL,
0.42 mmol, 2 equiv) via a syringe under N₂. The reaction
mixture was stirred at room temperature for 5 h. The
reaction mixture was poured into a mixture of EtOAc (200
20 mL) and saturated aqueous NaHCO₃ solution (100 mL). The
organic layer was separated, and the aqueous layer was
extracted with EtOAc (100 mLx3). The organic solution
was combined, dried over Na₂SO₄, filtered, and
concentrated under reduced pressure. The residue was
25 purified by flash column chromatography using EtOAc:
hexanes (1:1) as the eluent and dried overnight in vacuo
to give 10-chloro-acetyl-10-DAB as a colorless solid:
yield, 123 mg (93%). mp 231-233 °C dec; [α]_D²⁰ -66° (EtOAc,
c = 0.45); ¹H NMR (400 MHz, CDCl₃) δ 1.11(s, 3 H, Me17),
30 1.12(s, 3 H, Me16), 1.63(s, 1 H, 1-OH), 1.69(s, 3 H,
Me19), 1.89(ddd, J = 14.6, 10.9, 2.1 Hz, 1 H, H6b),
2.07(d, J = 5.2 Hz, 1 H, 13-OH), 2.09(d, J = 1.2, 3 H,
Me18), 2.12(d, J = 4.5 Hz, 1 H, 7-OH), 2.29(s, 3 H,
4-Ac), 2.30(m, 2 H, H14a, H14b), 2.58(ddd, J = 14.6, 9.7,
35 6.7 Hz, 1 H, H6a), 3.88(d, J = 7.0 Hz, 1 H, H3), 4.16(d,

41

$J = 8.3$ Hz, 1 H, H20b), 4.27(br s, 2 H, ClCH₂), 4.31(d, $J = 8.3$ Hz, 1 H, H20a), 4.44(ddd, $J = 10.9, 6.7, 4.5$ Hz, 1 H, H7), 4.90(m, 1 H, H13), 4.98(dd, $J = 9.7, 2.1$ Hz, 1 H, H5), 5.64(d, $J = 7.0$ Hz, 1 H, H2), 6.41(s, 1 H, H10),
 5 7.49(dd, $J = 7.9, 7.4$ Hz, 2 H, benzoate, *m*), 7.61(tt, $J = 7.4, 1.3$ Hz, 1 H, benzoate, *p*), 8.11(d, $J = 7.9, 1.3$ Hz, 2 H, benzoate, *o*) ppm. ¹³C NMR (75 MHz, CDCl₃) δ
 9.3(Me(19)), 15.3(Me(18)), 20.6(4-Ac), 22.3, 26.7(Me16, Me17), 35.8(C(6)), 38.6(C(14)), 40.5(ClCH₂), 42.6(C(15)),
 10 46.2(C(3)), 58.8(C(8)), 68.0(C(13)), 72.0, 75.0,
 75.9(C(7), C(2), C(10), C(20)), 79.0(C(1)), 80.9(C(4)),
 84.4(C(5)), 128.8, 129.7, 130.3, 133.9(benzoate),
 131.8(C(11)), 147.1(C(12)), 167.4, 167.7(ClCH₂C(O)O, benzoate), 171.0(4-Ac), 203.7(C(9)) ppm. Anal. Calcd for
 15 C₃₁H₃₇ClO₁₁. H₂O: C, 58.26; H, 6.15. Found: C, 58.26; H, 6.07.

10-Propionyl-10-DAB. To a solution of 10-DAB (47 mg, 0.086 mmol) in THF (2 mL) at room temperature was added a mixture of propionic anhydride anhydride (4 mL)
 20 and ZnCl₂/ THF solution (0.5 M, 350 mL, 0.173 mmol, 2 equiv) under N₂. The reaction mixture was stirred at room temperature for 14 h. Then the reaction mixture was diluted with EtOAc (150 mL), exhaustively washed with saturated aqueous NaHCO₃ solution (50 mLx3), brine. The
 25 organic layer was dried over Na₂SO₄, concentrated under reduced pressure. The residue was purified by flash column chromatography using EtOAc: hexanes (1:1) as the eluent and dried in vacuo to give 10-propionyl-10-DAB as a white solid: yield, 48 mg (93%). mp 212-213 °C dec;
 30 $[\alpha]_{\text{Hg}} -96^\circ$ (CHCl₃, $c = 0.78$); ¹H NMR (400 MHz, CDCl₃) δ
 1.11(s, 6 H, Me16, Me17), 1.24(t, $J = 7.6$ Hz, 3 H, CH₃CH₂), 1.60(s, 1 H, 1-OH), 1.67(s, 3 H, Me19), 1.87(ddd, $J = 14.7, 10.9, 2.2$ Hz, 1 H, H6b), 2.05(d, $J = 5.1$ Hz, 1 H, 13-OH), 2.06(d, $J = 1.3$ Hz, 3 H, Me18), 2.28(s, 3 H,
 35 4-Ac), 2.30(d, $J = 7.5$ Hz, 2 H, H14a, H14b), 2.51(d, $J =$

4.1 Hz, 1 H, 7-OH), 2.55(q, J = 7.6 Hz, 2 H, CH₃CH₂),
2.57(ddd, J = 14.7, 9.5, 6.7 Hz, 1 H, H6a), 3.90(d, J =
6.9 Hz, 1 H, H3), 4.16(dd, J = 8.4, 0.8 Hz, 1 H, H20b),
4.31(d, J = 8.4 Hz, 1 H, H20a), 4.48(ddd, J = 10.9, 6.7,
5 4.1 Hz, 1 H, H7), 4.90(m, 1 H, H13), 4.99(dd, J = 9.5,
2.2 Hz, 1 H, H5), 5.63(d, J = 6.9 Hz, 1 H, H2), 6.34(s, 1
H, H10), 7.48(dd, J = 8.1, 7.4 Hz, 2 H, benzoate, m),
7.61(tt, J = 7.4, 1.3 Hz, 1 H, benzoate, p), 8.11(dd, J =
8.3, 1.3 Hz, 2 H, benzoate, o) ppm. ¹³C NMR (75 MHz,
10 CDCl₃) δ 8.8(CH₃CH₂), 9.2(Me(19)), 15.2(Me(18)),
20.7(4-Ac), 22.3, 26.8, 27.4(Me16, Me17, CH₃CH₂),
35.5(C(6)), 38.7(C(14)), 42.6(C(15)), 46.1(C(3)),
58.7(C(8)), 67.9(C(13)), 72.3, 75.1, 76.1, 76.5(C(7),
C(2), C(10), C(20)), 79.1(C(1)), 80.9(C(4)), 84.5(C(5)),
15 128.7, 129.7, 130.3, 133.8(benzoate), 132.3(C(11)),
146.5(C(12)), 167.4(benzoate), 170.9, 174.9(4-Ac,
10-C(O)O), 204.6(C(9)) ppm. Anal. Calcd for C₃₂H₄₀O₁₁: C,
63.99; H, 6.71. Found: C, 63.81; H, 6.80.

20 C. Selective acylation of a C(10) hydroxyl group using
CeCl₃:

General procedure: To a solution of 10-DAB in THF
(20 mL per mmol of 10-DAB) under N₂ was added CeCl₃ and
the appropriate anhydride or pyrocarbonate (amounts
specified in Table 1). The reaction mixture was stirred
25 at 25 °C and monitored by TLC analysis. When that
analysis indicated complete reaction (time specified in
Table 1), the reaction mixture was diluted with EtOAc and
washed three times with saturated aqueous sodium
bicarbonate solution. The combined bicarbonate washings
30 were extracted three times with EtOAc, the organic layers
were combined and dried over sodium sulfate, and the
solvent was evaporated. The crude product was purified
by flash column chromatography. Further purification, if
necessary, was carried out by recrystallization from
35 EtOAc/Hexane.

Table 1. CeCl₃ catalyzed acylation of 10-DAB

$10\text{-DAB} \xrightarrow[\text{THF, RT}]{(\text{RCO})_2\text{O, CeCl}_3} 10\text{-Acyl-10-DAB}$				
Entry	R (eq)	CeCl ₃ (eq)	Time (hr)	Yield (%)
1	Me (10)	0.1	1.5	91
2	Pr (10)	0.1	3	100
3	iPr (10)	0.1	4.5	100
4	Ph (10)	0.1	21	94
5	cyclopropyl (10)	0.1	20.5	94
6	MeCH=CH (10)	0.1	20	91
7	CH ₂ =CHCH ₂ O (5)	0.1	1	96
8	EtO (5)	0.1	3	99
9	MeO (5)	0.1	3	98
10	tBuO (10)	0.7	24	94
11	BnO (3)	0.7	1	98

10-butyl-10-DAB. mp 145-149 °C; $[\alpha]_{\text{D}}^{25} -86.6'$ (CHCl₃, c = 1); ¹H NMR (500 MHz, CDCl₃) δ 8.13-8.11 (2H, m), 7.62 (1H, m), 7.51-7.48 (2H, m), 6.35 (1H, s), 5.64 (1H, d, J 7.0Hz), 4.99 (1H, d, J 7.7Hz), 4.90 (1H, m), 4.48 (1H, m), 4.31 (1H, d, J 8.3Hz), 4.18 (1H, d, J 8.3Hz), 3.91 (1H, d, J 7.0Hz), 2.60-2.42 (4H, m), 2.36-2.26 (2H, m), 2.28 (3H, s), 2.06 (3H, d, J 1.0Hz), 1.88 (1H, ddd, J 1.9, 10.9, 13.0Hz), 1.76 (2H, hex, J 7.4Hz), 1.68 (3H, s), 1.12 (6H, s) and 1.04 (3H, t, J 7.4Hz); ¹³C NMR (100MHz, CDCl₃) δ 204.2, 173.9, 170.6, 167.1, 146.2, 133.7, 132.0, 130.1, 129.4, 128.6, 84.5, 88.9, 79.1, 76.5, 76.0, 75.0, 72.3, 68.0, 58.8, 46.2, 42.7, 38.7, 37.1, 36.2, 35.6, 30.6, 27.0, 22.6, 20.9, 18.4, 17.8, 15.5 and 9.4; Anal. Calcd. for C₃₃H₄₂O₁₁: C, 64.48; H, 6.89 Found: C, 63.67; H, 7.01.

10-isobutyryl-10-DAB. mp 143 °C; $[\alpha]_{\text{Hg}} -62.6^\circ$
(CHCl₃, c=0.075); ¹H NMR (CDCl₃, 500MHz): δ 8.12 (2H, d, *J* 7.3Hz), 7.62 (1H, m), 7.51-7.48 (2H, m), 6.33 (1H, s), 5.65 (1H, d, *J* 7.3Hz), 5.00 (1H, d, *J* 7.9Hz), 4.91 (1H, m), 4.48 (1H, ddd, *J* 4.3, 6.7, 11.0Hz), 4.31 (1H, d, *J* 8.6Hz), 4.18 (1H, d, *J* 8.6Hz), 3.91 (1H, d, *J* 7.3Hz), 2.74 (1H, pent, *J* 6.7Hz), 2.57 (1H, m), 2.51 (1H, d, *J* 4.3Hz), 2.31 (1H, m), 2.28 (3H, s), 2.06 (3H, s), 2.01 (1H, d, *J* 5.5Hz), 1.90 (1H, ddd, *J* 2.3, 11.0, 14.6Hz), 1.68 (3H, s), 1.60 (1H, s), 1.51 (3H, s), 1.33 (3H, d, *J* 6.7Hz), 1.26 (3H, d, *J* 6.7Hz), 1.13 (3H, s) and 1.12 (3H, s); ¹³C NMR (100MHz, CDCl₃) δ 204.1, 177.2, 170.6, 167.1, 146.2, 133.7, 132.1, 130.1, 129.4, 128.6, 95.5, 84.5, 80.9, 79.1, 76.5, 75.8, 74.9, 72.3, 68.0, 58.8, 46.2, 42.7, 38.7, 35.6, 34.1, 27.0, 22.6, 20.9, 19.2, 18.7, 15.5 and 9.4; Anal. Calcd. for C₃₃H₄₂O₁₁•0.5H₂O: C, 64.48; H, 6.89 Found: C, 63.05; H, 6.70.

10-benzoyl-10-DAB. ¹H NMR (CDCl₃, 500MHz): δ 8.15-8.11 (4H, m), 7.64-7.6 (2H, m), 7.52-7.48 (4H, m), 6.62 (1H, s), 5.7 (1H, d, *J* 7.1Hz), 5.02 (1H, d, *J* 7.7Hz), 4.94 (1H, m), 4.57 (1H, ddd, *J* 4.4, 7.1, 11.0Hz), 4.33 (1H, d, *J* 8.2Hz), 4.20 (1H, d, *J* 8.3Hz), 3.99 (1H, d, *J* 6.6Hz), 2.62 (1H, ddd, *J* 6.6, 9.3, 14.8), 2.55 (1H, d, *J* 4.4Hz), 2.35 (2H, m), 2.30 (3H, s), 2.13 (3H, d, *J* 1.1Hz), 2.03 (1H, d, *J* 4.9Hz), 1.91 (1H, ddd, *J* 2.2, 11.0, 13.2Hz), 1.71 (3H, s), 1.65 (1H, s), 1.25 (3H, s) and 1.21 (3H, s); ¹³C NMR (100MHz, CDCl₃) δ 204.0, 170.7, 167.1, 166.5, 146.5, 133.7, 133.6, 132.0, 130.1, 129.9, 129.4, 129.3, 128.7, 128.5, 84.5, 80.9, 79.1, 76.5, 75.0, 72.4, 68.1, 58.8, 46.3, 42.8, 38.7, 35.8, 29.7, 27.2, 22.6, 21.2, 15.6 and 9.5; Anal. Calcd. for C₃₅H₄₀O₁₁: C, 66.66; H, 6.22 Found C, 66.46; H, 6.19.

45

10-*trans* crotonyl-10-DAB. ^1H NMR (CDCl_3 , 500MHz): δ
8.13 (2H, d, J 7.1Hz), 7.62 (1H, m), 7.51-7.48 (2H, m),
7.11 (1H, m), 6.42 (1H, s), 6.02 (1H, dq, J 1.7, 15.4Hz),
5.66 (1H, d, J 7.1Hz), 4.99 (1H, dd, J 2.0, 9.6Hz), 4.91
5 (1H, t, J 7.6Hz), 4.50 (1H, dd, J 7.1, 10.8Hz), 4.31 (1H,
d, J 8.3Hz), 4.19 (1H, d, J 8.3Hz), 3.93 (1H, d, J
7.1Hz), 2.61-2.55 (2H, m), 2.33-2.31 (2H, m), 2.28 (3H,
s), 2.07 (3H, d, J 1.5Hz), 1.95 (3H, dd, J 1.6, 6.8Hz),
1.89 (1H, ddd, J 2.3, 11.0, 13.4Hz), 1.68 (3H, s), 1.15
10 (3H, s) and 1.14 (3H, s); ^{13}C NMR (75MHz, CDCl_3) δ 212.4,
181.0, 170.8, 167.3, 166.5, 146.4, 133.8, 132.3, 130.2,
129.5, 128.7, 121.9, 116.0, 84.7, 84.6, 80.9, 79.2, 77.2,
75.9, 75.1, 72.4, 68.1, 58.8, 46.1, 42.7, 38.6, 35.6,
27.0, 20.9, 18.0, 15.4 and 9.3; Anal. Calcd. for $\text{C}_{33}\text{H}_{40}\text{O}_{11}$:
15 C, 64.69; H, 6.58 Found C, 63.93; H, 6.61.

10-cyclopropanoyl-10-DAB. ^1H (CDCl_3 , 500MHz): δ
8.12 (2H, d, J 7.3Hz), 7.62 (1H, t, J 7.5Hz), 7.49 (2H,
t, J 7.7Hz), 6.35 (1H, s), 5.65 (1H, d, J 7.0Hz), 4.99
(1H, app-d, J 8.2Hz), 4.91 (1H, m), 4.46 (1H, ddd, J 4.1,
20 6.8, 10.8Hz), 4.31 (1H, d, J 8.1Hz), 4.18 (1H, d, J
8.1Hz), 3.90 (1H, d, J 7.0Hz), 2.56 (1H, m), 2.51 (1H, d,
 J 4.1Hz), 2.31 (2H, m), 2.07 (3H, d, J 1.0Hz), 2.00 (1H,
d, J 4.9Hz), 1.87 (1H, ddd, J 2.1, 10.8, 14.6Hz), 1.79
(1H, ddd, J 3.4, 7.9, 12.4Hz), 1.68 (3H, s), 1.60 (1H,
25 s), 1.16-1.14 (2H, m), 1.13 (6H, s) and 1.01-0.97 (2H,
m); ^{13}C NMR (100MHz, CDCl_3) δ 204.3, 175.2, 170.6, 167.1,
146.4, 133.7, 132.0, 130.1, 129.4, 128.6, 84.5, 80.9,
79.1, 76.5, 76.0, 74.9, 72.4, 68.0, 58.8, 46.2, 42.7,
38.6, 35.6, 34.0, 27.0, 25.6, 24.9, 22.6, 21.0, 15.6,
30 13.1, 9.4 and 9.1; Anal. Calcd. for $\text{C}_{33}\text{H}_{40}\text{O}_{11}$: C, 64.69; H,
6.58 Found: C, 64.47; H, 6.66.

10-Ethoxycarbonyl-10-DAB. mp 214-215 °C; $[\alpha]_{\text{Hg}} -81^\circ$
(CHCl₃, c = 0.35); ¹H NMR (500 MHz, CDCl₃) δ 1.13(s, 3 H,
Me17), 1.14(s, 3 H, Me16), 1.38(t, J = 7.1 Hz, 3 H,
CH₃CH₂), 1.59(s, 1 H, 1-OH), 1.70(s, 3 H, Me19), 1.88(ddd,
5 J = 14.6, 10.5, 2.1 Hz, 1 H, H6b), 2.00(d, J = 5.0 Hz, 1
H, 13-OH), 2.10(d, J = 1.4 Hz, 3 H, Me18), 2.28(s, 3 H,
4-Ac), 2.30(m, 2 H, H14a, H14b), 2.46(d, J = 4.2 Hz, 1 H,
7-OH), 2.57(ddd, J = 14.6, 9.6, 6.7 Hz, 1 H, H6a),
3.88(d, J = 6.9 Hz, 1 H, H3), 4.18(d, J = 8.2 Hz, 1 H,
10 H20b), 4.31(d, J = 8.2 Hz, 1 H, H20a), 4.23-4.33(m, 2 H,
CH₃CH₂), 4.44(ddd, J = 10.5, 6.7, 4.2 Hz, 1 H, H7),
4.90(m, 1 H, H13), 4.98(dd, J = 9.6, 2.1 Hz, 1 H, H5),
5.65(d, J = 6.9 Hz, 1 H, H2), 6.17(s, 1 H, H10), 7.48(dd,
J = 8.2, 7.3 Hz, 2 H, benzoate, m), 7.60(tt, J = 7.3, 1.4
15 Hz, 1 H, benzoate, p), 8.11(d, J = 8.2, 1.4 Hz, 2 H,
benzoate, o) ppm; ¹³C NMR (75 MHz, CDCl₃) δ 9.2, 14.0,
15.5, 20.8, 22.4, 26.7, 35.4, 38.5, 42.4, 46.0, 58.6,
65.0, 67.7, 72.2, 74.9, 76.4, 78.7, 79.0, 80.6, 84.4,
128.7, 129.4, 130.1, 131.5, 133.7, 147.5, 155.4, 167.1,
20 170.8, 204.7 ppm.

10-Methoxycarbonyl-10-DAB. mp 218-219 °C; $[\alpha]_{\text{Hg}} -83^\circ$
(CHCl₃, c = 0.58); ¹H NMR (500 MHz, CDCl₃) δ 1.12(s, 3 H,
Me17), 1.13(s, 3 H, Me16), 1.59(s, 1 H, 1-OH), 1.70(s, 3
H, Me19), 1.88(ddd, J = 14.7, 10.8, 1.8 Hz, 1 H, H6b),
25 2.00(d, J = 5.0 Hz, 1 H, 13-OH), 2.10(d, J = 1.4 Hz, 3 H,
Me18), 2.28(s, 3 H, 4-Ac), 2.30(m, 2 H, H14a, H14b),
2.40(d, J = 4.1 Hz, 1 H, 7-OH), 2.57(ddd, J = 14.7, 9.7,
6.6 Hz, 1 H, H6a), 3.87(d, J = 6.9 Hz, 1 H, H3), 3.88(s,
3 H, MeOC(O)), 4.18(d, J = 8.4 Hz, 1 H, H20b), 4.31(d, J
30 = 8.4 Hz, 1 H, H20a), 4.44(ddd, J = 10.8, 6.6, 4.1 Hz, 1
H, H7), 4.90(m, 1 H, H13), 4.98(dd, J = 9.7, 1.8 Hz, 1 H,
H5), 5.65(d, J = 6.9 Hz, 1 H, H2), 6.17(s, 1 H, H10),
7.48(t, J = 8.2, 7.3 Hz, 2 H, benzoate, m), 7.61(tt, J =
7.3, 1.4 Hz, 1 H, benzoate, p), 8.11(d, J = 8.2, 1.4 Hz,
35 2 H, benzoate, o) ppm; ¹³C NMR (75 MHz, CDCl₃) δ 9.2,

47

15.5, 20.7, 22.4, 26.7, 35.5, 38.5, 42.4, 46.0, 55.4,
58.6, 65.0, 67.7, 72.1, 74.8, 76.4, 78.9, 79.0, 80.6,
84.4, 128.7, 129.4, 130.1, 131.4, 133.7, 147.5, 155.9,
167.1, 170.8, 204.6 ppm.

5 10-tBoc-10-DAB. mp 193-194 °C; $[\alpha]_{\text{D}}^{25}$ -82° (CHCl₃, c
=0.33); ¹H NMR (500 MHz, CDCl₃) δ 1.13(s, 6 H, Me17,
Me16), 1.48(s, 9 H, tBuO), 1.58(s, 1 H, 1-OH), 1.69(s, 3
H, Me19), 1.88(ddd, J = 14.9, 11.0, 2.2 Hz, 1 H, H6b),
1.99(d, J = 5.0 Hz, 1 H, 13-OH), 2.08(d, J = 1.4 Hz, 3 H,
10 Me18), 2.28(s, 3 H, 4-Ac), 2.30(m, 2 H, H14a, H14b),
2.56(ddd, J = 14.9, 9.6, 6.9 Hz, 1 H, H6a), 2.68(d, J =
3.6 Hz, 1 H, 7-OH), 3.88(d, J = 6.9 Hz, 1 H, H3), 4.19(d,
J = 8.2 Hz, 1 H, H20b), 4.31(d, J = 8.2 Hz, 1 H, H20a),
4.46(ddd, J = 11.0, 6.9, 3.6 Hz, 1 H, H7), 4.90(m, 1 H,
15 H13), 4.99(dd, J = 9.6, 2.2 Hz, 1 H, H5), 5.64(d, J = 6.9
Hz, 1 H, H2), 6.11(s, 1 H, H10), 7.48(t, J = 7.8 Hz, 2 H,
benzoate, m), 7.60(tt, J = 7.8, 1.3 Hz, 1 H, benzoate,
p), 8.11(dd, J = 7.8, 1.3 Hz, 2 H, benzoate, o) ppm; ¹³C
NMR (75 MHz, CDCl₃) δ 9.2, 15.6, 20.9, 22.4, 26.8, 27.5,
20 35.3, 38.5, 42.5, 45.9, 58.7, 67.9, 72.3, 74.7, 76.4,
78.0, 79.2, 80.8, 83.8, 84.5, 128.7, 129.4, 130.1, 131.8,
133.7, 147.3, 154.0, 167.2, 170.8, 205.0 ppm.

D. Selective carbamoylation of a C(10) hydroxyl group:

General procedure for the Selective Carbamoylation
25 of the C-10 Hydroxyl group of 10-DAB: A solution of
0.061 mmol (1.1 mol equiv) of the isocyanate in 2 mL of
THF was added, under nitrogen, to a mixture of 10-DAB (30
mg, 0.055 mmol) and CuCl (5.5 mg, 0.055 mmol) at 0 °C.
The mixture was stirred for the time indicated in Table
30 2. After this time the reaction was warmed to 25 °C and
stirring was continued for the time indicated in Table 2.
The reaction was quenched by the addition of saturated
aqueous ammonium chloride solution, and the mixture was
extracted three times with EtOAc. The combined organic

layers were washed with saturated aqueous NaHCO_3 solution, dried over sodium sulfate, and the solvent was evaporated to yield a white solid. The product was purified by flash column chromatography using 2:1 EtOAc/hexane as the eluent.

Table 2. Carbamoylation of 10-DAB

10-DAB $\xrightarrow{\text{RNCO, CuCl, THF}}$ 10-Carbamoyl-10-DAB				
Entry	R (eq)	Temp ($^{\circ}\text{C}$)	Time (hr)	Yield (%)
1	Et (1.1)	0	7.5	88
		rt	0.5	
2	allyl (1.1)	0	6	88
		rt	0.5	
3	Bu (1.1)	0	6.5	87
		rt	0.5	
4	Ph (1.1)	rt	3	94

10-ethylcarbamoyl-10-DAB. mp 241-243 $^{\circ}\text{C}$; $[\alpha]_{\text{D}}^{25} -92.0^{\circ}$ (CHCl_3 , $c=0.5$); ^1H NMR (400MHz, CDCl_3) δ 8.13 (2H, d, J 7.1Hz), 7.63 (1H, m), 7.52-7.48 (2H, m), 6.27 (1H, s), 5.63 (1H, d, J 6.9Hz), 5.01 (1H, dd, J 1.9, 9.6Hz), 4.97 (1H, m), 4.91 (1H, m), 4.50 (1H, ddd, J 3.7, 6.5, 10.5Hz), 4.31 (1H, d, J 8.3Hz), 4.17 (1H, d, J 8.3Hz), 3.88 (1H, d, J 7.0Hz), 3.32- 3.25 (2H, m), 3.10 (1H, d, J 3.7Hz), 2.56 (1H, ddd, J 6.8, 9.8, 14.8Hz), 2.31 (1H, m), 2.29 (3H, s), 2.09 (3H, s), 1.88 (1H, ddd, J 2.2; 11.0, 13.3Hz), 1.67 (3H, s), 1.60 (1H, s), 1.19 (3H, t, J 7.2Hz) and 1.10 (6H, s); Anal. Calcd. for $\text{C}_{32}\text{H}_{40}\text{NO}_{11}$: C, 62.43; H, 6.71 Found: C, 61.90; H, 6.77.

10-butylcarbamoyl-10-DAB. $[\alpha]_{\text{D}}^{25} -89.6^{\circ}$ (CHCl_3 , $c=0.25$); ^1H NMR (500MHz, CDCl_3) δ 8.12 (2H, d, J 7.3Hz), 7.61 (1H, m), 7.51- 7.45 (2H, m), 6.27 (1H, s), 5.64 (1H, d, J 6.7Hz), 5.00 (1H, d, J 8.0Hz), 4.91 (1H, m), 4.49

49

(1H, m), 4.31 (1H, d, J 8.5Hz), 4.19 (1H, d, J 8.5Hz), 3.89 (1H, d, J 6.7Hz), 3.25-3.23 (2H, m), 3.04 (1H, m), 2.56 (1H, ddd, J 6.7, 9.7, 14.7Hz), 2.30 (1H, d, J 7.9Hz), 2.28 (3H, s), 2.09 (3H, s), 1.99 (1H, d, J 4.9Hz), 1.88 (1H, ddd, J 2.5, 11.0, 13.4Hz), 1.68 (3H, s), 1.59 (1H, s), 1.55 (2H, b), 1.42-1.37 (2H, m), 1.11 (6H, s) and 0.95 (3H, t, J 7.6Hz); Anal. Calcd. for $C_{34}H_{44}NO_{11}$: C, 63.44; H, 7.05 Found: C, 62.64; H, 7.01.

10-phenylcarbamoyl-10-DAB. mp 178-180 °C; $[\alpha]_{\text{Hg}}$ -93.0° (CHCl_3 , $c=0.5$); ^1H NMR(400Hz, CDCl_3) δ 8.13 (2H, d, J 6.9Hz), 7.63 (1H, t, J 7.4Hz), 7.51 (2H, t, J 7.6Hz), 7.42 (1H, d, J 7.8Hz), 7.36-7.32 (2H, m), 7.12 (1H, t, J 7.4Hz), 6.87 (1H, b), 6.38 (1H, s), 5.66 (1H, d, J 7.0Hz), 5.02 (1H, app d, J 7.8Hz), 5.93 (1H, m), 4.52 (1H, ddd, J 3.8, 6.5, 10.5Hz), 4.33 (1H, d, J 8.3Hz), 4.18 (1H, d, J 8.3Hz), 3.91 (1H, d, J 7.0Hz), 2.83 (1H, d, J 4.0Hz), 2.59 (1H, ddd, J 6.5, 9.4, 14.5Hz), 2.33 (1H, m), 2.29 (3H, s), 2.12 (3H, d, J 1.4Hz), 2.04 (1H, d, J 5.1Hz), 1.89 (1H, ddd, J 2.2, 11.0, 14.4Hz), 1.69 (3H, s), 1.62 (1H, s), 1.15 (3H, s) and 1.13 (3H, s).

10-allylcarbamoyl-10-DAB. mp 165-170 °C; $[\alpha]_{\text{Hg}}$ -80.0° (CHCl_3 , $c=0.25$); ^1H NMR(500MHz, CDCl_3) δ 8.12 (2H, d, J 7.3Hz), 7.62 (1H, m), 7.51-7.48 (2H, m), 6.27 (1H, s), 5.89 (1H, m), 5.62 (1H, d, J 6.7Hz), 5.31 (1H, s), 5.19 (1H, d, J 9.8Hz), 5.08 (1H, m), 5.00 (1H, d, J 7.9Hz), 4.90 (1H, m), 4.49 (1H, ddd, J 3.7, 6.1, 10.4Hz), 4.31 (1H, d, J 8.5Hz), 4.17 (1H, d, J 8.5Hz), 3.88-3.86 (2H, m), 3.03 (1H, d, J 3.7Hz), 2.55 (1H, ddd, J 6.7, 9.8, 15.9Hz), 2.30 (1H, m), 2.29 (3H, s), 2.08 (3H, s), 2.06 (1H, app d, J 4.9Hz), 1.87 (1H, ddd, J 1.8, 11.0, 14.0Hz), 1.67 (3H, s), 1.58 (1H, s) and 1.09 (6H, s); Anal. Calcd. for $C_{33}H_{40}NO_{11}$: C, 63.15; H, 6.58 Found: C, 61.73; H, 6.45.

E. Selective silylation of a C(10) hydroxyl group:

10-TMS-10-DAB. To a solution of 10-DAB (100 mg, 0.18 mmol) in THF (10 mL) at 0 °C was slowly added N, O-bis(trimethylsilyl)trifluoroacetamide (1.0 mL, 3.7 mmol, 20 equiv) under N₂. The reaction mixture was stirred at 0 °C for 5 h. EtOAc (20 mL) was added, and the solution was filtered through a short column of silica gel. The silica gel was washed with EtOAc (100 mL), and the solution was concentrated under reduced pressure.

The residue was purified by flash column chromatography using EtOAc: hexanes (1:1) as the eluent and dried in vacuo overnight to give 10-TMS-10-DAB as a white solid: yield, 103 mg (91%). mp 189-191 °C; $[\alpha]_{\text{D}}^{25}$ -70° (CHCl₃, c = 0.55); ¹H NMR (400 MHz, CDCl₃) δ 0.18(s, 9 H, Me₃Si), 1.06(s, 3 H, Me17), 1.16(s, 3 H, Me16), 1.31(d, J = 8.6 Hz, 1 H, 7-OH), 1.56(s, 1 H, 1-OH), 1.68(s, 3 H, Me19), 1.79(ddd, J = 14.4, 11.1, 2.1 Hz, 1 H, H6b), 1.97(d, J = 4.9 Hz, 1 H, 13-OH), 2.03(d, J = 1.3 Hz, 3 H, Me18), 2.27(m, 2 H, H14a, H14b), 2.28(s, 3 H, 4-Ac), 2.58(ddd, J = 14.4, 9.6, 7.5 Hz, 1 H, H6a), 4.01(d, J = 7.2 Hz, 1 H, H3), 4.16(d, J = 8.2 Hz, 1 H, H20b), 4.25(ddd, J = 11.1, 8.6, 7.5 Hz, 1 H, H7), 4.30(d, J = 8.2 Hz, 1 H, H20a), 4.84(m, 1 H, H13), 4.97(dd, J = 9.6, 2.1 Hz, 1 H, H5), 5.27(s, 1 H, H10), 5.64(d, J = 7.2 Hz, 1 H, H2), 7.47(dd, J = 8.2, 7.5 Hz, 2 H, benzoate, m), 7.60(tt, J = 7.5, 1.2 Hz, 1 H, benzoate, p), 8.11(dd, J = 8.2, 1.2 Hz, 2 H, benzoate, o) ppm. ¹³C NMR (75 MHz, CDCl₃) δ 0.2(Me₃S), 9.7(Me(19)), 14.4(Me(18)), 19.6(4-Ac), 22.4, 26.6(Me16, Me17), 37.1(C(6)), 38.6(C(14)), 42.6(C(15)), 47.2(C(3)), 57.8(C(8)), 68.0(C(13)), 72.0, 75.1, 76.1, 76.8(C(7), C(2), C(10), C(20)), 78.9(C(1)), 81.2(C(4)), 84.3(C(5)), 128.8, 130.3, 133.7(benzoate), 137.0(C(11)), 139.0(C(12)), 167.4(benzoate), 171.0(4-Ac), 209.5(C(9)) ppm. Anal. Calcd for C₃₂H₄₄O₁₀Si. 1/2H₂O: C, 61.42; H, 7.25. Found: C, 61.61; H, 7.12.

10-TES-10-DAB. To a solution of 10-DAB (85 mg, 0.16 mmol) in THF (3 mL) at 0 °C was slowly added N, O-bis(triethylsilyl)trifluoroacetamide (484 mL, 1.56 mmol, 10 equiv), and a catalytic amount of LiHMDS/THF solution (1 M, 5 mL, 0.005 mmol), respectively, under N₂. The reaction mixture was stirred at 0 °C for 5 min. EtOAc (10 mL) was added, and the solution was filtered through a short column of silica gel. The silica gel was washed with EtOAc (100 mL), and the solution was concentrated under reduced pressure. The residue was purified by flash column chromatography using EtOAc: hexanes (1:2) as the eluent and dried *in vacuo* overnight to give the 10-TES-10-DAB as a white solid: yield, 98 mg (95 %). mp 234-235 °C dec; $[\alpha]_{\text{D}}^{25}$ -69° (CHCl₃, c = 0.95); IR 3690, 2958, 1714, 1602 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 0.68(m, 6 H, (CH₃CH₂)₃Si), 1.00(t, J = 7.9, 9 H, (CH₃CH₂)₃Si), 1.08(s, 3 H, Me17), 1.19(s, 3 H, Me16), 1.29(d, J = 8.4 Hz, 1 H, 7-OH), 1.55(s, 1 H, 1-OH), 1.69(s, 3 H, Me19), 1.79(ddd, J = 14.4, 11.0, 2.0 Hz, 1 H, H6b), 1.92(d, J = 5.0 Hz, 1 H, 13-OH), 2.03(d, J = 1.0 Hz, 3 H, Me18), 2.27(s, 3 H, 4-Ac), 2.29(m, 2 H, H14a, H14b), 2.59(ddd, J = 14.4, 9.5, 6.7 Hz, 1 H, H6a), 4.02(d, J = 7.2 Hz, 1 H, H3), 4.18(d, J = 8.5 Hz, 1 H, H20b), 4.23(ddd, J = 11.0, 8.4, 6.7 Hz, 1 H, H7), 4.30(d, J = 8.5 Hz, 1 H, H20a), 4.86(m, 1 H, H13), 4.97(dd, J = 9.5, 2.0 Hz, 1 H, H5), 5.28(s, 1 H, H10), 5.66(d, J = 7.2 Hz, 1 H, H2), 7.47(dd, J = 7.9, 7.9 Hz, 2 H, benzoate, m), 7.59(tt, J = 7.9, 1.0 Hz, 1 H, benzoate, p), 8.11(dd, J = 7.9, 1.0 Hz, 2 H, benzoate, o) ppm. ¹³C NMR (75 MHz, CDCl₃) δ 4.9, 6.5(TES), 9.7(Me(19)), 14.3(Me(18)), 19.6(4-Ac), 22.4, 26.6(Me16, Me17), 37.1(C(6)), 38.6(C(14)), 42.6(C(15)), 47.3(C(3)), 57.9(C(8)), 67.9(C(13)), 71.9, 75.1, 76.1, 76.7(C(7), C(2), C(10), C(20)), 78.9(C(1)), 81.2(C(4)), 84.3(C(5)),

128.7, 129.9, 130.3, 133.7(benzoate), 137.0(C(11)),
138.8(C(12)), 167.4(benzoate), 171.0(4-Ac), 209.5(C(9))
ppm. Anal. Calcd for $C_{35}H_{50}O_{10}Si$. H_2O : C, 62.11; H, 7.74.
Found: C, 62.45; H, 7.74.

5

Example 2

General Procedure for the Preparation of 7-silyl-10-TES-10-DAB. To a solution of 7-triethylsilyl-10-DAB, 7-t-butylldimethylsilyl-10-DAB, or 7-dimethylisopropylsilyl-10-DAB in THF at 0 °C was slowly added N, O-bis(triethylsilyl)trifluoroacetamide (5 equiv), and a catalytic amount of LiHMDS/THF solution (5 mol %), respectively, under N_2 . The reaction mixture was stirred at 0 °C for 15 min. EtOAc (10 mL) was added, and the solution was filtered through a short column of silica gel. The silica gel was washed with EtOAc (100 mL), and the solution was concentrated under reduced pressure. The residue was purified by flash column chromatography using EtOAc: hexanes (1:2) as the eluent and dried *in vacuo* overnight to give, respectively, 7,10-bis(triethylsilyl)-10-DAB (95% yield), 7-t-butylldimethylsilyl-10-triethylsilyl-10-DAB (98% yield), or 7-dimethylisopropylsilyl-10-triethylsilyl-10-DAB (94% yield).

7-Dimethylphenylsilyl-10-TBS-10-DAB. To a solution of 7-dimethylphenylsilyl-10-DAB (35 mg, 0.052 mmol) in THF (2 mL) at 0 °C was added N,O-bis(t-butylldimethylsilyl)trifluoroacetamide (337 μ L, 1.09 mmol, 20 equiv), and a catalytic amount of LiHMDS/THF solution (1 M, 6 μ L, 0.006 mmol), respectively, under N_2 . The reaction mixture was stirred at 0 °C for 4 h, then warmed to room temperature for an additional 4 h. EtOAc (10 mL) was added, and the solution was filtered through a short column of silica gel. The silica gel was washed with EtOAc (100 mL), and the solution was concentrated under reduced pressure. The residue was purified by flash

column chromatography using EtOAc: hexanes (1:2) as the eluent and dried in vacuo overnight to give 39 mg (92% yield) of 7-dimethylphenylsilyl-10-*t*-butyldimethylsilyl-10-DAB.

5

Example 3**Selective silylation of a C(7) hydroxyl group**

7-TBS-10-DAB. To a mixture of 10-DAB (38 mg, 0.070 mmol), imidazole (190 mg, 2.79 mmol, 40 equiv), and *tert*-butyldimethylsilyl chloride (210 mg, 1.40 mmol, 20 equiv) was added DMF (0.1 mL) at room temperature under N₂. The reaction mixture was vigorously stirred at room temperature for 24 h. EtOAc (20 mL) was added, and the solution was filtered through a short column of silica gel. The silica gel was washed with EtOAc (200 mL), and the solution was concentrated under reduced pressure. The residue was purified by flash column chromatography using 10% EtOAc-CH₂Cl₂ as the eluent and dried in vacuo overnight to give 7-TBS-10-DAB as a white solid: yield, 41 mg (90%). mp 222-223 °C; [α]_D²⁰ -51° (CHCl₃, c = 0.36); ¹H NMR (400 MHz, CDCl₃) δ 0.05, 0.06 (2 s, 6 H, Me₂Si), 0.83 (s, 9 H, Me₃C), 1.09 (s, 6 H, Me₁₆, Me₁₇), 1.57 (s, 1 H, 1-OH), 1.75 (s, 3 H, Me₁₉), 1.87 (ddd, J = 14.4, 10.6, 2.0 Hz, 1 H, H_{6b}), 2.01 (d, J = 5.0 Hz, 1 H, 13-OH), 2.09 (d, J = 1.3, 3 H, Me₁₈), 2.28 (m, 2 H, H_{14a}, H_{14b}), 2.29 (s, 3 H, 4-Ac), 2.46 (ddd, J = 14.4, 9.6, 6.7 Hz, 1 H, H_{6a}), 3.96 (d, J = 6.9 Hz, 1 H, H₃), 4.16 (d, J = 8.3 Hz, 1 H, H_{20b}), 4.24 (d, J = 2.2 Hz, 1 H, 10-OH), 4.31 (d, J = 8.3 Hz, 1 H, H_{20a}), 4.38 (dd, J = 10.6, 6.7 Hz, 1 H, H₇), 4.88 (m, 1 H, H₁₃), 4.96 (dd, J = 9.6, 2.0 Hz, 1 H, H₅), 5.15 (d, J = 2.0 Hz, 1 H, H₁₀), 5.60 (d, J = 6.9 Hz, 1 H, H₂), 7.47 (dd, J = 8.1, 7.5 Hz, 2 H, benzoate, m), 7.60 (tt, J = 7.5, 1.3 Hz, 1 H, benzoate, p), 8.10 (d, J = 8.1, 1.3 Hz, 2 H, benzoate, o) ppm. ¹³C NMR (75 MHz, CDCl₃) δ -5.8, -3.8 (Me₂Si), 9.7 (Me₁₉), 14.8 (Me₁₈), 17.6 (Me₃C), 19.3 (4-Ac), 22.4, 26.7 (Me₁₆, Me₁₇),

54

25.4 (Me₃C), 37.4 (C(6)), 38.7 (C(14)), 42.7 (C(15)),
 47.0 (C(3)), 58.0 (C(8)), 68.0 (C(13)), 73.1, 74.7,
 75.0 (C(7), C(2), C(10), C(20)), 78.9 (C(1)), 80.9 (C(4)),
 84.3 (C(5)), 128.8, 129.8, 130.3, 133.8 (benzoate),
 5 135.7 (C(11)), 141.9 (C(12)), 167.4 (benzoate), 171.2 (4-Ac),
 210.8 (C(9)) ppm. Anal. Calcd for C₃₅H₅₀O₁₀Si: C, 63.80; H,
 7.65. Found: C, 63.72; H, 7.70.

7-Dimethylphenylsilyl-10-DAB. To a THF (3 mL)
 solution of 10-DAB (54 mg, 0.099 mmol) at -20 °C was added
 10 pyridine (0.6 mL), dimethylphenylsilyl chloride (250 mL,
 1.49 mmol, 15 equiv) under N₂. The reaction mixture was
 stirred at -20 °C for 2 h. EtOAc (10 mL) and saturated
 NaHCO₃ aqueous solution (0.5 mL) was added, and the
 solution was quickly filtered through a short column of
 15 silica gel. The silica gel was washed with EtOAc (100
 mL), and the solution was concentrated under reduced
 pressure. The residue was purified by flash column
 chromatography using EtOAc: CH₂Cl₂ (1:10) as the eluent
 and dried overnight *in vacuo* to give 7-dimethylphenyl-
 20 silyl-10-DAB as a white solid: yield, 62 mg (92%). mp
 219-220 °C; [α]_D²⁰ -28° (CHCl₃, c = 0.27); ¹H NMR (400 MHz,
 CDCl₃) δ 0.35, 0.37 (2 s, 6 H, Me₂Si), 1.05 (s, 3 H, Me17),
 1.06 (s, 3 H, Me16), 1.54 (s, 1 H, 1-OH), 1.73 (d, J = 1.1,
 3 H, Me18), 1.76 (s, 3 H, Me19), 1.90 (ddd, J = 14.4, 10.6,
 25 2.1 Hz, 1 H, H6b), 1.93 (d, J = 5.0 Hz, 1 H, 13-OH),
 2.23 (m, 2 H, H14a, H14b), 2.25 (s, 3 H, 4-Ac), 2.43 (ddd, J
 = 14.4, 9.6, 6.8 Hz, 1 H, H6a), 3.86 (d, J = 7.0 Hz, 1 H,
 H3), 4.10 (d, J = 2.1 Hz, 1 H, 10-OH), 4.16 (d, J = 8.3 Hz,
 1 H, H20b), 4.28 (d, J = 8.3 Hz, 1 H, H20a), 4.31 (dd, J =
 30 10.6, 6.8 Hz, 1 H, H7), 4.81 (m, 1 H, H13), 4.84 (d, J =
 2.1 Hz, 1 H, H10), 4.90 (dd, J = 9.6, 2.1 Hz, 1 H, H5),
 5.59 (d, J = 7.0 Hz, 1 H, H2), 7.41, 7.53 (2 m, 5 H, C₆H₅),
 7.46 (dd, J = 8.0, 7.5 Hz, 2 H, benzoate, m), 7.55 (tt, J =
 7.5, 1.2 Hz, 1 H, benzoate, p), 8.09 (d, J = 8.0, 1.2 Hz,
 35 2 H, benzoate, o) ppm. ¹³C NMR (75 MHz, CDCl₃) δ -1.8,

55

-1.1 (Me₂Si), 9.8 (Me(19)), 14.4 (Me(18)), 19.4 (4-Ac), 22.3, 26.7 (Me16, Me17), 37.2 (C(6)), 38.6 (C(14)), 42.6 (C(15)), 46.7 (C(3)), 58.0 (C(8)), 68.0 (C(13)), 73.2, 74.7, 75.0 (C(7), C(2), C(10), C(20)), 78.8 (C(1)), 80.8 (C(4)), 84.3 (C(5)), 128.3, 128.8, 129.8, 130.2, 130.3, 133.65, 133.74 (PhSi, benzoate), 135.4 (C(11)), 142.1 (C(12)), 167.4 (benzoate), 171.0 (4-Ac), 210.9 (C(9)) ppm. Anal. Calcd for C₃₇H₄₆O₁₀Si. 1/2H₂O: C, 64.61; H, 6.89. Found: C, 64.72; H, 6.81.

10 7-Dimethylisopropylsilyl-10-DAB. To a solution of
10-DAB (97 mg, 0.18 mmol) in pyridine (1 mL) at -10 °C was
added dimethylisopropylsilyl chloride (580 mL, 3.57 mmol,
20 equiv) under N₂. The reaction mixture was stirred at
-10 °C for 3 h. EtOAc (10 mL) was added, and the solution
15 was quickly filtered through a short column of silica
gel. The silica gel was washed with EtOAc (150 mL), and
the solution was concentrated under reduced pressure.
The residue was purified by flash column chromatography
using EtOAc: hexanes (1:2) as the eluent to give
20 7-dimethylisopropyl-10-DAB as a white solid: yield, 107
mg (93%). mp 229-230 °C; [α]_D²⁰ -56° (CHCl₃, c = 0.62); ¹H
NMR (400 MHz, CDCl₃) δ 0.05, 0.06 (2 s, 6 H, Me₂Si),
0.70 (m, 1 H, CHSi), 0.90, 0.92 (2 dd, J = 7.4, 1.7, 6 H,
Me₂CH), 1.09 (s, 6 H, Me16, Me17), 1.56 (s, 1 H, 1-OH),
25 1.74 (s, 3 H, Me19), 1.89 (ddd, J = 14.4, 10.6, 2.1 Hz, 1
H, H6b), 1.99 (d, J = 5.0 Hz, 1 H, 13-OH), 2.09 (d, J =
1.4, 3 H, Me18), 2.28 (d, J = 7.9, 2 H, H14a, H14b),
2.29 (s, 3 H, 4-Ac), 2.44 (ddd, J = 14.4, 9.7, 6.7 Hz, 1 H,
H6a), 3.96 (d, J = 7.3 Hz, 1 H, H3), 4.17 (d, J = 8.3 Hz, 1
30 H, H20b), 4.24 (d, J = 2.2 Hz, 1 H, 10-OH), 4.31 (d, J =
8.3 Hz, 1 H, H20a), 4.38 (dd, J = 10.6, 6.7 Hz, 1 H, H7),
4.85 (m, 1 H, H13), 4.95 (dd, J = 9.7, 2.1 Hz, 1 H, H5),
5.15 (d, J = 2.2 Hz, 1 H, H10), 5.61 (d, J = 7.3 Hz, 1 H,
H2), 7.47 (dd, J = 8.2, 7.5 Hz, 2 H, benzoate, m),
35 7.60 (tt, J = 7.5, 1.4 Hz, 1 H, benzoate, p), 8.10 (d, J =

56

8.2, 1.4 Hz, 2 H, benzoate, o) ppm. ^{13}C NMR (75 MHz, CDCl_3) δ -4.6, -3.3 (Me_2Si), 9.7 (Me(19)), 14.8, 14.9 (CHSi, Me(18)), 16.4, 16.5 (Me_2CH), 19.4 (4-Ac), 22.4, 26.7 (Me16, Me17), 37.3 (C(6)), 38.7 (C(14)), 42.7 (C(15)), 47.0 (C(3)), 58.0 (C(8)), 68.0 (C(13)), 73.1, 74.7, 75.0 (C(7), C(2), C(10), C(20)), 78.9 (C(1)), 80.9 (C(4)), 84.3 (C(5)), 128.8, 129.8, 130.3, 133.7 (benzoate), 135.7 (C(11)), 142.0 (C(12)), 167.4 (benzoate), 171.1 (4-Ac), 210.8 (C(9)) ppm. Anal. Calcd for $\text{C}_{34}\text{H}_{48}\text{O}_{10}\text{Si}$. H_2O : C, 61.61; H, 7.60. Found: C, 61.30; H, 7.35.

7-Tribenzylsilyl-10-DAB. To a mixture of 10-DAB (62 mg, 0.11 mmol), imidazole (280 mg, 4.11 mmol, 36 equiv) and tribenzylsilyl chloride (364 mg, 1.14 mmol, 10 equiv) was added DMF (0.4 mL) under N_2 . The reaction mixture was stirred at room temperature for 3 h. EtOAc (30 mL) was added, and the solution was filtered through a short column of silica gel. The silica gel was washed with EtOAc (150 mL), and the solution was concentrated under reduced pressure. The residue was purified twice by flash column chromatography, first time using EtOAc: hexanes (1:2) as the eluent, second time using EtOAc: CH_2Cl_2 as the eluent, and dried overnight in vacuo to give the 7-tribenzylsilyl-10-DAB as a white solid: yield, 88 mg (91%). mp 161-163 $^\circ\text{C}$; IR 3690, 2928, 2890, 1712, 1600 cm^{-1} ; $[\alpha]_{\text{Hg}} -46^\circ$ (CHCl_3 , $c = 0.46$); ^1H NMR (400 MHz, CDCl_3) δ 1.10 (s, 3 H, Me17, Me16), 1.56 (s, 1 H, 1-OH), 1.71 (ddd, $J = 14.2, 10.9, 2.0$ Hz, 1 H, H6b), 1.74 (s, 3 H, Me19), 2.00 (d, $J = 5.1$ Hz, 1 H, 13-OH), 2.07 (ddd, $J = 14.2, 9.6, 6.6$ Hz, 1 H, H6a), 2.10 (d, $J = 1.2$, 3 H, Me18), 2.12 (s, 6 H, $(\text{PhCH}_2)_3\text{Si}$), 2.27 (d, $J = 7.5$ Hz, 2 H, H14a, H14b), 2.27 (s, 3 H, 4-Ac), 3.99 (d, $J = 7.0$ Hz, 1 H, H3), 4.16 (d, $J = 8.5$ Hz, 1 H, H20b), 4.18 (d, $J = 2.2$ Hz, 1 H, 10-OH), 4.28 (d, $J = 8.5$ Hz, 1 H, H20a), 4.58 (dd, $J = 10.9, 6.6$ Hz, 1 H, H7), 4.81 (dd, $J = 9.6, 2.0$ Hz, 1 H, H5), 4.89 (m, 1 H, H13), 5.21 (d, $J = 2.2$ Hz, 1 H, H10), 5.61 (d, $J = 7.0$

57

Hz, 1 H, H₂), 6.93, 7.09, 7.20 (3 m, 15 H, (PhCH₂)₃Si),
7.48 (dd, J = 8.1, 7.5 Hz, 2 H, benzoate, m), 7.61 (tt, J =
7.5, 1.3 Hz, 1 H, benzoate, p), 8.10 (d, J = 8.1, 1.3 Hz,
2 H, benzoate, o) ppm. ¹³C NMR (75 MHz, CDCl₃) δ
5 9.9 (Me(19)), 15.0 (Me(18)), 19.5 (4-Ac), 22.4, 26.7 (Me16,
Me17), 23.6 (Si(CH₂Ph)₃), 36.9 (C(6)), 38.7 (C(14)),
42.7 (C(15)), 46.8 (C(3)), 58.0 (C(8)), 68.0 (C(13)), 74.4,
74.9, 75.0 (C(7), C(2), C(10), C(20)), 78.8 (C(1)),
80.8 (C(4)), 84.1 (C(5)), 124.9, 128.7, 128.8, 129.1,
10 129.8, 130.3, 133.8, 137.8 (Si(CH₂Ph)₃, benzoate),
135.5 (C(11)), 142.2 (C(12)), 167.4 (benzoate), 170.9 (4-Ac),
210.8 (C(9)) ppm. Anal. Calcd for C₅₀H₅₆O₁₀Si. 1/2H₂O: C,
70.32; H, 6.73. Found: C, 70.11; H, 6.57.

Example 4

15 Selective acylation of 10-acyl-10-DAB

10-Alloc-7-p-Nitrobenzyloxycarbonyl-10-DAB. To a
mixture of 10-alloc-10-DAB (33 mg, 0.053 mmol) and DMAP
(19.3 mg, 0.16 mmol, 3 equiv) in dichloromethane (4 mL)
at 0 °C was added a dichloromethane solution (1 mL) of
20 p-nitrobenzyl chloroformate (23 mL, 0.11 mmol, 2 equiv)
under N₂. The reaction mixture was stirred at 0 °C for 4
h. EtOAc (10 mL) was added, the solution was quickly
filtered through a short column of silica gel. The
silica gel was washed with EtOAc (100 mL), and the
25 solution was concentrated under reduced pressure. The
residue was purified by flash column chromatography using
EtOAc: hexanes (1:2) as the eluent and dried overnight in
vacuo to give 10-alloc-7-p-nitrobenzyloxycarbonyl-10-DAB
as a colorless solid: yield, 34 mg (92%).

30 7-Benzyloxycarbonyl baccatin III. To a stirred
solution of baccatin III (100 mg, 0.168 mmol) in
methylene chloride under N₂ at room temperature was added
4-dimethylaminopyridine (204 mg, 1.68 mmol) followed by
addition of benzyl chloroformate (240 mL, 1.68 mmol).

The reaction mixture was stirred at room temperature and the progress of the reaction was monitored by TLC. After about 4 h the reaction was complete. The mixture was diluted with EtOAc (10 mL) and was transferred to a separatory funnel containing 50 mL of a 50% EtOAc/Hexanes. The mixture was washed with saturated sodium bicarbonate and the organic layer was separated. The aqueous layer was washed with 20 mL of 50% EtOAc/Hexanes. The combined organic layers were washed with brine, dried over MgSO_4 , and concentrated under reduced pressure. The crude product was passed through a short column to give 115 mg (95%) of a white solid m.p. 245-248°C; $[\alpha]_D^{25}$ -60.5° c (C=0.007, CHCl_3). ^1H NMR (CDCl_3 , 400 MHz) δ 8.10 (d, J=9.6 Hz, 2H, o-benzoate), 7.60-6.8 (m, 8H, benzoate, Bn), 6.45 (s, 1H, H10), 5.63 (d, J=6.9 Hz, 1H, H2b), 5.56 (dd, J=10.6, 7.2 Hz, 1H, H7), 5.56 (dd, J=18.5, 12.0 Hz, 2H, Bn), 4.97 (d, J=10.6, 1H, H5), 4.87 (m, 1H, H13), 4.31 (d, J=10.5, 1H, H20a), 4.15 (d, J=10.5, 1H, H20b), 4.02 (d, J=6.9, 1H, H3), 2.61 (m, 1H, H6a), 2.30 (m, 2H, H14's), 2.29 (s, 3H, 4Ac), 2.18 (s, 3H, 10Ac), 2.15 (br s, 3H, Me18), 2.08 (d, J=5.2 Hz, 13OH), 1.94 (m, 1H, 6b), 1.79 (s, 3H, Me19), 1.58 (s, 1H, 10H), 1.14 (s, 3H, Me16), 1.09 (s, 3H, Me17).

7-Allyloxycarbonyl baccatin III. To a stirred solution of baccatin III (30 mg, 0.051 mmol) in methylene chloride (1 mL) under N_2 at room temperature, was added 4-dimethylaminopyridine (62.3 mg, 0.51 mmol) followed by addition of allyl chloroformate (54 mL, 0.51 mmol). The reaction mixture was stirred at room temperature and the progress of the reaction was followed by TLC. After about 1.5 h the reaction was complete. The mixture was diluted by EtOAc (5 mL) and was transferred to a separatory funnel containing 50 mL of a 50% EtOAc/Hexanes. The mixture was washed with saturated sodium bicarbonate and the organic layer was separated.

The aqueous layer was washed with 10 mL of 50% EtOAc/Hexanes, the combined organic layers were washed with brine, dried over MgSO_4 , and concentrated under reduced pressure. The crude product was passed through a short column to give 33.1 mg (97%) of a white solid m.p. 239-244 °C; $[\alpha]^{25}_D$ -61.5 c (0.01, CHCl_3) ^1H NMR (CDCl_3 , 500 MHz) δ 8.12(d, $J=8.3$ Hz, 2H, o--benzoate), 7.66-7.45 (m, 3H, benzoate), 6.43(s, 1H, H10), 5.97(m, 1H, int. allyl), 5.64(d, $J=7.0$ Hz, 1H, H2b), 5.54 (dd, $J=10.5, 7.0$ Hz, 1H, H7), 5.28(m, 2H, ext. allyl), 4.97(d, $J=9.6$ Hz, 1H, H5), 4.87(m, 1H, H13), 4.67(m, 2H, CH_2 allyl), 4.31(d, $J=8.5$ Hz, 1H, H20a), 4.17(d, $J=8.5$, 1H, H20b), 4.02(d, $J=7.0$, 1H, H3), 2.64(m, 1H, H6a), 2.30(d, $J=8.0$ Hz, 2H, H14's), 2.29(s, 3H, 4Ac), 2.16(s, 3H, 10Ac), 2.15(br s, 3H, Me18), 2.01(d, $J=5$ Hz, 13OH), 1.96(m, 1H, 6b), 1.81(s, 3H, Me19), 1.58(s, 1H, 10H), 1.15(s, 3H, Me16), 1.02(s, 3H, Me17).

Example 5

Selective ketalization of 10-acyl-10-DAB

7-MOP baccatin III. To a solution of baccatin III (101 mg, 0.172 mmol) in THF (8 mL) at -20 °C was added 2-methoxypropene (0.66 mL, 6.89 mmol, 40 equiv), followed by the addition of a catalytic amount of toluenesulfonic acid (0.1 M solution in THF, 43 μL , 0.004 mmol, 0.025 equiv) under N_2 . The reaction mixture was stirred at -20 °C for 3 h. TLC analysis indicated complete consumption of the starting material and the formation of desired product as the only major product. Triethylamine (0.5 mL) was added, and the solution was warmed to room temperature, diluted with EtOAc (100 mL), washed with saturated aqueous NaHCO_3 solution, dried over Na_2SO_4 , and concentrated under reduced pressure. The residue was dried *in vacuo* overnight to give 112 mg (99%) of crude product. Recrystallization of the crude product from EtOAc/hexanes gave 105 mg (93%) of 7-MOP baccatin III as

a white crystal, mp 181-183 °C; ¹H NMR (500 MHz, C₆D₆) δ 1.01 (s, 3 H, Me17), 1.11 (br s, 1 H, 13-OH), 1.28 (s, 3 H, Me16), 1.39, 1.78 (2 s, 6 H, Me₂CO), 1.62 (s, 1 H, 1-OH), 1.78 (s, 3 H, 10-Ac), 1.92 (s, 3 H, 4-Ac), 2.09 (s, 3 H, Me18), 2.12 (s, 3 H, Me19), 2.14 (ddd, J = 15.0, 10.9, 2.2 Hz, 1 H, H6b), 2.18 (dd, J = 15.6, 9.4 Hz, 1 H, H14b), 2.31 (dd, J = 15.6, 7.0 Hz, 1 H, H14b), 2.97 (s, 3 H, MeO), 3.15 (ddd, J = 15.0, 9.9, 6.7 Hz, 1 H, H6a), 4.08 (d, J = 7.0 Hz, 1 H, H3), 4.24 (m, 1 H, H13), 4.33 (d, J = 8.3 Hz, 1 H, H20b), 4.41 (d, J = 8.3 Hz, 1 H, H20a), 4.78 (dd, J = 10.9, 6.7 Hz, 1 H, H7), 4.97 (dd, J = 9.9, 2.2 Hz, 1 H, H5), 5.95 (d, J = 7.0 Hz, 1 H, H2), 6.79 (s, 1 H, H10), 7.15 (m, 3 H, benzoate, m, p), 8.28 (d, J = 8.0 Hz, 2 H, benzoate, o) ppm; Anal. Calcd for C₃₅H₄₆O₁₂: C, 63.82; H, 7.04. Found: C, 63.72; H, 7.07.

Example 6

Selective acylation of 10-silyl-10-DAB

7-Acetyl-10-TES-10-DAB. To a stirred solution of 10-TES-10-DAB (65 mg, 0.098 mmol) in dichloromethane (4 ml) at 0 °C under N₂, was added DMAP (36 mg, 0.296 mmol, 3 equiv), followed by addition of acetic anhydride (14 mL, 0.148 mmol, 1.5 equiv). The reaction mixture was stirred at 0 °C for 4.5 hrs and the TLC analysis indicated the complete consumption of starting material. The reaction mixture was then filtered through a short pad of silica gel, the silica gel was washed with EtOAc (100 mL) and the solution was concentrated under reduced pressure. The residue was purified by flash chromatography using EtOAc:hexanes (1:2) as the eluent and dried in vacuo overnight to give the 7-acetyl-10-TES-10-DAB: yield, 65.7 mg (95%). ¹H NMR (CDCl₃, 400 MHz), δ 0.60 (m, 6 H, (CH₃CH₂)₃Si), 0.97 (t, J = 7.9 Hz, 9 H, (CH₃CH₂)₃Si), 1.05 (s, 3 H, Me17), 1.18 (s, 3 H, Me16), 1.56 (s, 1 H, 1-OH), 1.79 (s, 3 H, Me19), 1.83 (ddd, J = 14.5, 10.3, 2.0 Hz, 1 H, H6b), 1.97 (m, 1 H, 13-OH), 2.00 (s, 3 H, 7-Ac), 2.07 (d,

61

$J = 1.3$ Hz, 3 H, Me18), 2.26(m, 2 H, H14a, H14b), 2.29(s, 3 H, 4-Ac), 2.57(ddd, $J = 14.5, 9.5, 7.3$ Hz, 1 H, H6a), 4.06(d, $J = 7.0$ Hz, 1 H, H3), 4.17(d, $J = 8.2$ Hz, 1 H, H20b), 4.31(d, $J = 8.2$ Hz, 1 H, H20a), 4.84(m, 1 H, H13),
 5 4.94(dd, $J = 9.5, 2.0$ Hz, 1 H, H5), 5.29(s, 1 H, H10), 5.46(dd, $J = 10.3, 7.3$ Hz, 1 H, H7), 5.65(d, $J = 7.0$ Hz, 1 H, H2), 7.47(m, 2 H, benzoate, m), 7.60(m, 1 H, benzoate, p), 8.11(d, $J = 8.0$ Hz, 2 H, benzoate, o) ppm.

7-Troc-10-TES-10-DAB. To a mixture of 10-TES-10-DAB
 10 (40 mg, 0.061 mmol) and DMAP (72 mg, 0.61 mmol, 10 equiv) in dichloromethane (2 mL) was added trichloroethyl chloroformate (24 mL, 0.184 mmol, 3 equiv) under N_2 . The reaction mixture was stirred at room temperature and the progress of the reaction was monitored by TLC analysis.
 15 After 0.5 h, TLC analysis indicated almost complete disappearance of 10-TES-10-DAB and the formation of the product as the only major spot. Methanol (5 mL) was added, and the solution was quickly filtered through a short column of silica gel. The silica gel was washed
 20 with EtOAc (100 mL), and the solution was concentrated under reduced pressure. The residue was purified by flash chromatography using EtOAc: CH_2Cl_2 (1:10) as the eluent and dried in vacuo overnight to give 7-Troc-10-TES- 10-DAB as a white solid: yield, 49 mg
 25 (97%); 1H NMR (500 MHz, $CDCl_3$) δ 0.61(m, 6 H, $(CH_3CH_2)_3Si$), 0.99(t, $J = 7.9, 9$ Hz, $(CH_3CH_2)_3Si$), 1.08(s, 3 H, Me17), 1.20(s, 3 H, Me16), 1.56(s, 1 H, 1-OH), 1.84(s, 3 H, Me19), 1.96(d, $J = 4.9$ Hz, 1 H, 13-OH), 2.01(ddd, $J = 14.4, 10.5, 2.0$ Hz, 1 H, H6b), 2.08(d, $J = 1.2, 3$ Hz, Me18), 2.29(m, 2 H, H14a, H14b), 2.29(s, 3 H, 4-Ac),
 30 2.68(ddd, $J = 14.4, 9.5, 7.3$ Hz, 1 H, H6a), 4.08(d, $J = 6.7$ Hz, 1 H, H3), 4.18(d, $J = 8.5$ Hz, 1 H, H20b), 4.32(d, $J = 8.5$ Hz, 1 H, H20a), 4.43(d, $J = 11.9$ Hz, 1 H, $CHH'OC(O)$), 4.86(m, 1 H, H13), 4.95(dd, $J = 9.3, 2.0$ Hz,
 35 1 H, H5), 4.98(d, $J = 11.9$ Hz, 1 H, $CHH'OC(O)$), 5.33(s, 1

H, H10), 5.37(dd, $J = 10.5, 7.3$ Hz, 1 H, H7), 5.67(d, $J = 6.7$ Hz, 1 H, H2), 7.48(dd, $J = 7.9, 7.3$ Hz, 2 H, benzoate, m), 7.60(tt, $J = 7.3, 1.2$ Hz, 1 H, benzoate, p), 8.11(dd, $J = 7.9, 1.2$ Hz, 2 H, benzoate, o) ppm.

5 7-*p*-Nitrobenzyloxycarbonyl-10-TES-10-DAB. To a mixture of 10-TES-10-DAB (40 mg, 0.061 mmol) and DMAP (72 mg, 0.61 mmol, 10 equiv) in dry chloroform (2 mL) was added *p*-nitrobenzyl chloroformate (131 mg, 0.61 mmol, 10 equiv) under N₂. The reaction mixture was stirred at room
10 temperature and the progress of the reaction was monitored by TLC analysis. After 45 min, TLC analysis indicated almost complete disappearance of 10-TES-10-DAB and the formation of the product as the only major spot. Methanol (10 mL) was added, and the solution was quickly
15 filtered through a short column of silica gel. The silica gel was washed with EtOAc (100 mL), and the solution was concentrated under reduced pressure. The residue was purified by flash chromatography using EtOAc: CH₂Cl₂ (1:10) as the eluent and dried *in vacuo* overnight
20 to give 7-*p*-Nitrobenzyloxycarbonyl-10-TES-10-DAB as a white solid: yield, 48.3 mg (95%); ¹H NMR (500 MHz, CDCl₃) δ 0.60(m, 6 H, (CH₃CH₂)₃Si), 0.95(t, $J = 7.9$, 9 H, (CH₃CH₂)₃Si), 1.08(s, 3 H, Me17), 1.19(s, 3 H, Me16), 1.55(s, 1 H, 1-OH), 1.83(s, 3 H, Me19), 1.93(ddd, $J =$
25 14.3, 10.4, 2.2 Hz, 1 H, H6b), 1.96(d, $J = 4.9$ Hz, 1 H, 13-OH), 2.09(d, $J = 1.2$, 3 H, Me18), 2.29(m, 2 H, H14a, H14b), 2.29(s, 3 H, 4-Ac), 2.65(ddd, $J = 14.3, 9.3, 7.3$ Hz, 1 H, H6a), 4.08(d, $J = 7.0$ Hz, 1 H, H3), 4.18(d, $J = 8.6$ Hz, 1 H, H20b), 4.31(d, $J = 8.6$ Hz, 1 H, H20a),
30 4.86(m, 1 H, H13), 4.95(dd, $J = 9.3, 2.2$ Hz, 1 H, H5), 5.06(d, $J = 13.4$ Hz, 1 H, CHH'OC(O)), 5.31(d, $J = 13.4$ Hz, 1 H, CHH'OC(O)), 5.33(s, 1 H, H10), 5.36(dd, $J = 10.4, 7.3$ Hz, 1 H, H7), 5.66(d, $J = 7.0$ Hz, 1 H, H2),

7.48(dd, $J = 7.4, 7.3$ Hz, 2 H, benzoate, *m*), 7.53(d, $J = 8.9$ Hz, 2 H, $\text{NO}_2\text{C}_6\text{H}_4$), 7.59(tt, $J = 7.3, 1.2$ Hz, 1 H, benzoate, *p*), 8.12(dd, $J = 7.4, 1.2$ Hz, 2 H, benzoate, *o*), 8.23(d, $J = 8.9$ Hz, 2 H, $\text{NO}_2\text{C}_6\text{H}_4$) ppm.

5 7-Cbz-10-TES-10-DAB. To a mixture of 10-TES-10-DAB
(40 mg, 0.061 mmol) and DMAP (440 mg, 3.64 mmol, 60
8equiv) in dry chloroform (2 mL) was slowly added four
equal aliquot of benzyl chloroformate (4x130 mL, 3.64
mmol, 60 equiv) via a syringe in a 10-min interval under N_2
10 during a period of 40 min. The reaction mixture was then
stirred at room temperature and the progress of the
reaction was monitored by TLC analysis. After 2 h, TLC
analysis indicated almost complete disappearance of
10-TES-10-DAB and the formation of the product as the
15 only major spot. Methanol (10 mL) was added, and the
solution was poured into ethyl acetate (100 mL), washed
with a saturated aqueous NaHCO_3 solution, H_2O , and brine.
The solution was dried over Na_2SO_4 , concentrated under
reduced pressure. The residue was purified by flash
20 chromatography using EtOAc: CH_2Cl_2 (1:10) as the eluent
and dried in vacuo overnight to give 7-CBz-10-TES-10-DAB
as a white solid: yield, 45 mg (93%); ^1H NMR (500 MHz,
 CDCl_3) δ 0.62(m, 6 H, $(\text{CH}_3\text{CH}_2)_3\text{Si}$), 0.97(t, $J = 7.9$, 9 H,
 $(\text{CH}_3\text{CH}_2)_3\text{Si}$), 1.07(s, 3 H, Me17), 1.20(s, 3 H, Me16),
25 1.55(s, 1 H, 1-OH), 1.81(s, 3 H, Me19), 1.91(ddd, $J =$
14.3, 10.5, 2.1 Hz, 1 H, H6b), 1.96(d, $J = 4.9$ Hz, 1 H,
13-OH), 2.10(d, $J = 1.2$, 3 H, Me18), 2.28(m, 2 H, H14a,
H14b), 2.28(s, 3 H, 4-Ac), 2.64(ddd, $J = 14.3, 9.5, 7.3$
Hz, 1 H, H6a), 4.08(d, $J = 7.0$ Hz, 1 H, H3), 4.17(d, $J =$
30 8.5 Hz, 1 H, H20b), 4.30(d, $J = 8.5$ Hz, 1 H, H20a),
4.86(m, 1 H, H13), 4.95(dd, $J = 9.5, 2.1$ Hz, 1 H, H5),
5.01(d, $J = 12.2$ Hz, 1 H, $\text{CHH}'\text{OC}(\text{O})$), 5.24(d, $J = 12.2$
Hz, 1 H, $\text{CHH}'\text{OC}(\text{O})$), 5.34(s, 1 H, H10), 5.37(dd, $J =$

10.5, 7.3 Hz, 1 H, H7), 5.65(d, J = 7.0 Hz, 1 H, H2),
 7.32-7.37(m, 5 H, PhCH₂O), 7.47(dd, J = 8.3, 7.3 Hz, 2 H,
 benzoate, m), 7.59(tt, J = 7.3, 1.3 Hz, 1 H, benzoate,
 p), 8.12(dd, J = 8.3, 1.3 Hz, 2 H, benzoate, o) ppm.

5

Example 7**Selective silylation of 10-acyl-10-DAB**

7-dimethylisopropylsilyl baccatin III. To a stirred
 solution of baccatin III (30 mg, 0.051 mmol) in pyridine
 (0.6 mL) at 0 °C under N₂, was added chlorodimethyl-
 10 isopropylsilane (160 uL, 1.02 mmol). The reaction
 mixture was stirred at that temperature and the progress
 of the reaction was monitored by TLC. After about 1.5 h
 the reaction was complete. Ethyl acetate (5 mL) was
 added and the solution was transferred to a separatory
 15 funnel containing 50 mL of a 50% EtOAc /Hexanes. The
 mixture was washed with saturated sodium bicarbonate and
 the organic layer was separated. The aqueous layer was
 extracted with 10 mL of 50% EtOAc/Hexanes and the
 combined organic layers were washed with saturated sodium
 20 chloride, dried over MgSO₄, concentrated under reduced
 pressure. The crude product was passed through a short
 silica gel column to give 33.9 mg (97%) of a white solid
 m.p. 204-207 °C; [α]_D²⁵ -58.6° c (0.009, CHCl₃). ¹HNMR
 (CDCl₃, 500MHz), δ 8.10(d, J=8.4Hz, 2H, o-benzoate), 7.60-7.20
 25 (m, 3H, benzoate), 6.4 (s, 1H, H10), 5.64(d, J=7.1Hz, 1H, H2b),
 4.95(d, J=4.9Hz, 1H, H5), 4.84(m, 1H, H13), 4.44(dd, J=10.4, 6.8
 Hz, 1H, H7), 4.30(d, J=8.3 Hz, 1H, H20a), 4.14 (d, J=8.3 Hz,
 1H, H20b), 4.15(d, J=7.2 Hz, 1H, H3), 2.49(m, 1H, H6a),
 2.23(m, 2H, H14's), 2.28(s, 3H, 4Ac), 2.18(br s, 3H, Me
 30 18), 2.17(s, 3H, 10Ac), 2.01(d, J=5.0 Hz, 13 OH), 1.86(m,
 1H, 6b), 1.69(s, 3H, Me19), 1.61(s, 1H, 1OH), 1.20(s,
 3H, Me16), 1.05(s, 3H, Me17), 0.87(d, J=7.1 Hz, 6H, i-pr),
 0.73(m, 1H, i-pr), 0.09(s, 6H, Me2Si).

7-dimethylphenylsilyl baccatin III. To a stirred solution of baccatin III (20 mg, 0.034 mmol) in THF (1.25 mL) at -10 °C under N₂, was added chlorodimethylphenylsilane (68 uL, 0.41 mmol), followed by addition of pyridine (250 mL, 3.1 mmol). The reaction mixture was stirred at that temperature and the progress of the reaction was monitored by TLC. After about one hour the reaction was complete. Ethyl acetate (5 mL) was added and the solution was transferred to a separatory funnel containing 30 mL of 50% EtOAc /Hexanes. The mixture was washed with saturated sodium bicarbonate and the organic layer was separated. The aqueous layer was extracted with 10 mL of 50% EtOAc/Hexanes and the combined organic layers were washed with saturated sodium chloride, dried over MgSO₄, concentrated under reduced pressure. The crude product was passed through a short silica gel column to give 24.1 mg (98%) of a white solid m.p. 210-213 °C; $[\alpha]_D^{25}$ -58.3.5° c (0.005, CHCl₃) ¹H NMR (CDCl₃, 500 MHz) δ 8.35(d, J=8.5 Hz, 2H, o-benzoate), 7.627.25(m, 8H, benzoate, phenyl), 6.42 (s, 1H, H10), 5.64 (d, J=6.9 Hz, 1H, H2b), 4.84 (m, 1H, H5), 4.81 (m, 1H, H13), 4.46 (dd, J=10.6, 6.9 Hz, 1H, H7), 4.21 (d, J=8.5 Hz, 1H, H20a), 4.14 (d, J=8.5 Hz, 1H, H20b), 3.85 (d, J=6.9 Hz, 1H, H3), 2.34 (m, 1H, H6a), 2.26 (d, J=8 Hz, 2H, H14's), 2.24 (s, 3H, 4Ac), 2.15 (s, 3H, 10Ac), 2.02 (br d, J=1 Hz, 3H, Me 18), 1.93 (d, J=5 Hz, 1H, 13OH), 1.77 (m, 1H, 6b), 1.72 (s, 3H, Me19), 1.59 (s, 1H, 1OH), 1.20 (s, 3H, Me16), 1.05 (s, 3H, Me17), 0.446 (s, 3H, Me Si), 0.335 (s, 3H, Me Si).

7-dimethylphenylsilyl-10-propionyl-10-DAB. To a stirred solution of 10-propionyl-10-DAB (0.200 g, 0.333 mmol) in THF (12 mL) at -10 °C, was added chlorodimethylphenylsilane (0.668 mL, 4.00 mmol) followed by pyridine dropwise (2.48 mL, 30.64 mmol). The reaction was stirred for 90 minutes. Ethyl acetate (20 mL) was added and the solution transferred to a separatory funnel containing

100 mL of 50% EtOAc/Hexanes. The mixture was washed with saturated sodium bicarbonate and the organic layer separated. The aqueous layer was extracted with 50%EtOAc/Hexanes (30 mL) and the combined organic
5 extracts washed with saturated sodium chloride, dried over Na₂SO₄, concentrated in vacuo. The crude solid was then purified with flash column chromatography using 50% EtOAc/hexane as eluent to give 7-dimethylphenylsilyl-10-propionyl-10-DAB (0.242 g, 99%) as a solid. ¹H NMR
10 (CDCl₃, 500 MHz), δ 0.34, 0.45(2 s, 6 H, Me₂Si), 1.05(s, 3 H, Me17), 1.20(t, J = 7.5 Hz, 3 H, CH₃CH₂), 1.21(s, 3 H, Me16), 1.60(s, 1 H, 1-OH), 1.72(s, 3 H, Me19), 1.78(ddd, J = 14.5, 10.0, 2.0 Hz, 1 H, H6b), 2.04(m, 1 H, 13-OH), 2.05(s, 3 H, Me18), 2.27(m, 2 H, H14a, H14b), 2.25(s, 3
15 H, 4-Ac), 2.34(ddd, J = 14.5, 9.5, 7.0 Hz, 1 H, H6a), 2.42, 2.49(2 dq, J = 16.5, 7.5 Hz, 6 H, CH₃CH₂), 3.87(d, J = 7.5 Hz, 1 H, H3), 4.14(d, J = 8.0 Hz, 1 H, H20b), 4.27(d, J = 8.0 Hz, 1 H, H20a), 4.47(dd, J = 10.0, 7.0
Hz, 1 H, H7), 4.82(m, 1 H, H13), 4.85(dd, J = 9.5, 2.0
20 Hz, 1 H, H5), 5.64(d, J = 7.5 Hz, 1 H, H2), 6.44(s, 1 H, H10), 7.32-7.36, 7.55-7.57(2 m, 5 H, PhSi), 7.46(m, 2 H, benzoate, m), 7.59(m, 1 H, benzoate, p), 8.10(d, J = 8.0
Hz, 2 H, benzoate, o) ppm.

7-Dimethylphenylsilyl-10-cyclopropanecarbonyl-
25 10-DAB. To a solution of 10-cyclopropanecarbonyl-10-DAB (680 mg, 1.1 mmol) in THF (25 mL) were added with stirring pyridine (3.5 mL) and then chlorodimethylphenylsilane (1.8 mL, 11 mmol) at -10 °C under N₂. The solution was stirred till the reaction completed. Then
30 quenched with sat. NaHCO₃ (20 mL). The mixture was extracted with EtOAc (2 x 250 mL). The combined organic layers were washed with brine (2 x 10 mL), dried and filtered. Concentration of the filtrate in vacuo and followed by flash chromatography (hexane:EtOAc, 4:1) gave
35 7-Dimethyl-phenylsilyl-10-cyclopropane-carbonyl-10-DAB

(816 mg, ~100%). ^1H NMR (CDCl_3 , 500 MHz), δ 0.32, 0.43 (2 s, 6 H, Me_2Si), 0.91, 1.00, 1.17 (3 m, 5 H, cyclopropyl), 1.07 (s, 3 H, Me17), 1.21 (s, 3 H, Me16), 1.73 (s, 3 H, Me19), 1.74 (s, 1 H, 1-OH), 1.78 (ddd, $J = 14.4, 10.5, 2.1$ Hz, 1 H, H6b), 2.04 (m, 1 H, 13-OH), 2.05 (d, $J = 1.5$ Hz, 3 H, Me18), 2.24 (s, 3 H, 4-Ac), 2.26 (m, 2 H, H14a, H14b), 2.34 (ddd, $J = 14.4, 9.5, 6.7$ Hz, 1 H, H6a), 3.87 (d, $J = 7.0$ Hz, 1 H, H3), 4.15 (d, $J = 8.2$ Hz, 1 H, H20b), 4.26 (d, $J = 8.2$ Hz, 1 H, H20a), 4.46 (dd, $J = 10.5, 6.7$ Hz, 1 H, H7), 4.82 (m, 1 H, H13), 4.85 (dd, $J = 9.5, 2.1$ Hz, 1 H, H5), 5.65 (d, $J = 7.0$ Hz, 1 H, H2), 6.44 (s, 1 H, H10), 7.32-7.36, 7.55-7.57 (2 m, 5 H, PhSi), 7.46 (m, 2 H, benzoate, m), 7.59 (m, 1 H, benzoate, p), 8.10 (d, $J = 8.0$ Hz, 2 H, benzoate, o) ppm.

15

Example 8

7-*p*-Nitrobenzyloxycarbonyl-10-DAB. To a THF solution (1 mL) of 10-alloc-7-*p*-nitrobenzyloxycarbonyl-10-DAB (34 mg, 0.048 mmol) at room temperature was added a THF solution (1 mL) of formic acid (19 mL, 0.48 mmol, 10 equiv) and butylamine (47 mL, 0.48 mmol, 10 equiv), followed by the addition of $\text{Pd}(\text{PPh}_3)_4$ under N_2 . The reaction mixture was stirred at room temperature for 0.5 h. EtOAc (10 mL) was added, and the solution was quickly filtered through a short column of silica gel. The silica gel was washed with EtOAc (100 mL), and the solution was concentrated under reduced pressure. The residue was purified by flash column chromatography using EtOAc: hexanes (1:2) as the eluent and dried *in vacuo* to give 7-*p*-nitrobenzyloxycarbonyl-10-DAB as a colorless solid: yield, 28 mg (93%). $[\alpha]_{\text{D}}^{25} -38^\circ$ (CHCl_3 , $c = 0.48$); ^1H NMR (400 MHz, CDCl_3) δ 1.06 (s, 3 H, Me16), 1.09 (s, 3 H, Me17), 1.55 (s, 1 H, 1-OH), 1.86 (s, 3 H, Me19), 2.01 (ddd, $J = 14.4, 10.7, 2.0$ Hz, 1 H, H6b), 2.03 (d, $J = 5.1$ Hz, 1 H, 13-OH), 2.09 (d, $J = 1.3, 3$ H, Me18), 2.28 (m, 2 H, H14a, H14b), 2.30 (s, 3 H, 4-Ac), 2.62 (ddd, $J = 14.4, 9.5,$

68

7.3 Hz, 1 H, H6a), 3.89(d, J = 2.0 Hz, 1 H, 10-OH),
 4.08(d, J = 6.9 Hz, 1 H, H3), 4.20(d, J = 8.4 Hz, 1 H,
 H20b), 4.34(d, J = 8.4 Hz, 1 H, H20a), 4.88(m, 1 H, H13),
 4.96(dd, J = 9.5, 2.0 Hz, 1 H, H5), 5.19(d, J = 13.3, 1
 5 H, CHH'OC(O)), 5.26(d, J = 13.3, 1 H, CHH'OC(O)),
 5.36(dd, J = 10.7, 7.3 Hz, 1 H, H7), 5.40(d, J = 2.0 Hz,
 1 H, H10), 5.64(d, J = 6.9 Hz, 1 H, H2), 7.48(dd, J =
 8.1, 7.5 Hz, 2 H, benzoate, m), 7.52(d, J = 8.7, 2 H,
 NO₂C₆H₄), 7.61(tt, J = 7.5, 1.3 Hz, 1 H, benzoate, p),
 10 8.10(d, J = 8.1, 1.3 Hz, 2 H, benzoate, o), 8.26(d, J =
 8.7, 2 H, NO₂C₆H₄) ppm. ¹³C NMR (75 MHz, CDCl₃) δ
 10.5(Me(19)), 14.6(Me(18)), 19.4(4-Ac), 22.2, 26.4(Me16,
 Me17), 33.2(C(6)), 38.7(C(14)), 42.4(C(15)), 46.5(C(3)),
 56.5(C(8)), 67.9, 68.3(C(13), OCH₂Ph-NO₂-p), 74.7, 75.2,
 15 76.8(C(7), C(2), C(10), C(20)), 78.8(C(1)), 80.4(C(4)),
 83.6(C(5)), 124.1, 128.4, 128.9, 130.3, 133.9
 (OCH₂Ph-NO₂-p, benzoate), 135.0(C(11)), 142.4, 143.0
 (OCH₂Ph-NO₂-p, C(12)), 154.2(OC(O)O), 167.3(benzoate),
 171.1(4-Ac), 211.6(C(9))ppm.

20

Example 9

Selective Esterification of the C-10 Hydroxyl of
10-DAB using the catalytic DyCl₃ reaction: A solution of
 butyric anhydride (0.55 mmol) in THF (1.32 ml) was added,
 under a nitrogen atmosphere, to a solid mixture of 10-DAB
 25 (30 mg, 0.055 mmol) and DyCl₃ (1.3 mg, 10 mol.% wrt
 10-DAB). The resulting suspension was stirred at room
 temperature until judged complete by TLC (2:1
 EtOAc/Hexane). The reaction was diluted with EtOAc and
 washed three times with saturated NaHCO₃ solution. The
 30 combined bicarbonate washings were extracted three times
 with EtOAc, these combined organics were dried (Na₂SO₄)
 and the solvent evaporated. The crude product was
 triturated with hexanes and the mother liquors decanted

away. Crystallization from EtOAc/hexanes yielded 10-butyl-10-DAB identical to that isolated from the CeCl_3 catalysed reaction.

Example 10


5 *Selective Esterification of the C-10 Hydroxyl of 10-DAB using the catalytic YbCl_3 reaction:* A solution of butyric anhydride (0.55 mmol) in THF (1.32 ml) was added, under a nitrogen atmosphere, to a solid mixture of 10-DAB (30 mg, 0.055 mmol) and YbCl_3 (1.3 mg, 10 mol.% wrt
10 10-DAB). The resulting suspension was stirred at room temperature until judged complete by TLC (2:1 EtOAc/Hexane). The reaction was diluted with EtOAc and washed three times with saturated NaHCO_3 solution. The combined bicarbonate washings were extracted three times
15 with EtOAc, these combined organics were dried (Na_2SO_4) and the solvent evaporated. The crude product was triturated with hexanes and the mother liquors decanted away. Crystallization from EtOAc/hexanes yielded 10-butyl-10-DAB identical to that isolated from the CeCl_3 ,
20 catalysed reaction.

In view of the above, it will be seen that the several objects of the invention are achieved.

As various changes could be made in the above compositions without departing from the scope of the
25 invention, it is intended that all matter contained in the above description be interpreted as illustrative and not in a limiting sense.

What is claimed is:


1. A process for the acylation of a C(10) hydroxy group of a taxane, the process comprising
treating the taxane with an acylating agent in a reaction mixture containing less than one equivalent of a
5 base for each equivalent of taxane to form a C(10) acylated taxane.
2. The process of claim 1 wherein the taxane has C(7) and C(10) hydroxy groups and the acylating agent selectively reacts with the C(10) hydroxy group.
3. The process of claim 1 wherein the taxane reacted with the acylating agent is 10-deacetyl baccatin III.
4. The process of claim 1 wherein the taxane has the structure:



wherein

- 5 R_1 is hydrogen, hydroxy, protected hydroxy, or together with R_{14} or R_2 forms a carbonate;
- R_2 is keto, $-OT_2$, acyloxy, or together with R_1 forms a carbonate;
- R_4 is $-OT_4$ or acyloxy;
- 10 R_7 is hydrogen, halogen, $-OT_7$, $-OCOZ_7$, or $-OCOOZ_7$;
- R_9 is hydrogen, keto, $-OT_9$, $-OCOZ_9$, or $-OCOOZ_9$;
- R_{10} is hydroxy;
- R_{13} is hydroxy, protected hydroxy, keto, or

71



- 15 R_{14} is hydrogen, $-OT_{14}$, acyloxy, or together with R_1
forms a carbonate;
 T_2 , T_4 , T_7 , T_9 and T_{14} are independently hydrogen or
hydroxy protecting group;
 X_1 is $-OX_6$, $-SX_7$, or $-NX_8X_9$;
20 X_2 is hydrogen, hydrocarbyl, substituted hydrocarbyl,
or heteroaryl;
 X_3 and X_4 are independently hydrogen, hydrocarbyl,
substituted hydrocarbyl, or heteroaryl;
 X_5 is $-X_{10}$, $-OX_{10}$, $-SX_{10}$, $-NX_8X_{10}$, or $-SO_2X_{11}$;
25 X_6 is hydrocarbyl, substituted hydrocarbyl,
heteroaryl, hydroxy protecting group or a functional
group which increases the water solubility of the taxane
derivative;
 X_7 is hydrocarbyl, substituted hydrocarbyl,
30 heteroaryl, or sulfhydryl protecting group;
 X_8 is hydrogen, hydrocarbyl, or substituted
hydrocarbyl;
 X_9 is an amino protecting group;
 X_{10} is hydrocarbyl, substituted hydrocarbyl, or
35 heteroaryl;
 X_{11} is hydrocarbyl, substituted hydrocarbyl,
heteroaryl, $-OX_{10}$, or $-NX_8X_{14}$; and
 X_{14} is hydrogen, hydrocarbyl, substituted
hydrocarbyl, or heteroaryl.

5. The process of claim 4 wherein

R₁ is hydroxy or together with R₁₄ or R₂ forms a carbonate;


5 R_2 is $-OCOZ_2$, $-OCOOZ_2$, or together with R_1 forms a carbonate;

72

R_4 is $-\text{OCOZ}_4$;

R_9 is hydrogen or keto;

R_{13} is hydroxy, protected hydroxy, or



- 10 R_{14} is hydrogen, hydroxy, protected hydroxy, or together with R_1 forms a carbonate;

X_1 is $-\text{OX}_6$ or $-\text{NX}_8\text{X}_9$;

X_2 is hydrogen, hydrocarbyl, or substituted hydrocarbyl;

- 15 X_3 and X_4 are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl;

X_5 is $-\text{X}_{10}$, $-\text{OX}_{10}$, or $-\text{NX}_8\text{X}_{10}$;

X_6 is a hydroxy protecting group;

- 20 X_8 is hydrogen, hydrocarbyl, or substituted hydrocarbyl;

X_9 is an amino protecting group;

X_{10} is hydrocarbyl, substituted hydrocarbyl, or heteroaryl; and

- 25 Z_2 and Z_4 are independently hydrocarbyl, substituted hydrocarbyl, or heteroaryl.

6. The process of claim 1 wherein the acylating agent is selected from the group consisting of anhydrides, dicarbonates, thiodicarbonates, and isocyanates.

7. The process of claim 6 wherein the reaction mixture contains a Lewis acid.


73

8. The process of claim 7 wherein the taxane reacted with the acylating agent is 10-deacetyl baccatin III.

9. The process of claim 1 wherein the reaction mixture contains a Lewis acid.

10. The process of claim 9 wherein the taxane reacted with the acylating agent is 10-deacetyl baccatin III.

11. The process of claim 9 wherein the taxane has the structure:



wherein

5 R_1 is hydrogen, hydroxy, protected hydroxy, or together with R_{14} or R_2 forms a carbonate;

R_2 is keto, $-OT_2$, acyloxy, or together with R_1 forms a carbonate;


R_4 is $-OT_4$ or acyloxy;

10 R_7 is hydrogen, halogen, $-OT_7$, $-OCOZ_7$, or $-OCOOZ_7$;

R_9 is hydrogen, keto, $-OT_9$, $-OCOZ_9$, or $-OCOOZ_9$;

R_{10} is hydroxy;

R_{13} is hydroxy, protected hydroxy, keto, or



15 R_{14} is hydrogen, $-OT_{14}$, acyloxy, or together with R_1 forms a carbonate;

T_2 , T_4 , T_7 , T_9 and T_{14} are independently hydrogen or hydroxy protecting group;

X_1 is $-OX_6$, $-SX_7$, or $-NX_8X_9$;

20 X_2 is hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl;

X_3 and X_4 are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl;

X_5 is $-X_{10}$, $-OX_{10}$, $-SX_{10}$, $-NX_8X_{10}$, or $-SO_2X_{11}$;

25 X_6 is hydrocarbyl, substituted hydrocarbyl, heteroaryl, hydroxy protecting group or a functional group which increases the water solubility of the taxane derivative;

30 X_7 is hydrocarbyl, substituted hydrocarbyl, heteroaryl, or sulfhydryl protecting group;

X_8 is hydrogen, hydrocarbyl, or substituted hydrocarbyl;

X_9 is an amino protecting group;

35 X_{10} is hydrocarbyl, substituted hydrocarbyl, or heteroaryl;

X_{11} is hydrocarbyl, substituted hydrocarbyl, heteroaryl, $-OX_{10}$, or $-NX_8X_{14}$; and

X_{14} is hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl.

12. The process of claim 11 wherein

R_1 is hydroxy or together with R_{14} or R_2 forms a carbonate;


5 R_2 is $-OCOZ_2$, $-OCOOZ_2$, or together with R_1 forms a carbonate;

R_4 is $-OCOZ_4$;

R_9 is hydrogen or keto;

R_{13} is hydroxy, protected hydroxy, or

75



- 10 R_{14} is hydrogen, hydroxy, protected hydroxy, or together with R_1 forms a carbonate;
 X_1 is $-OX_6$ or $-NX_8X_9$;
 X_2 is hydrogen, hydrocarbyl, or substituted hydrocarbyl;
- 15 X_3 and X_4 are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl;
 X_5 is $-X_{10}$, $-OX_{10}$, or $-NX_8X_{10}$;
 X_6 is a hydroxy protecting group;
 X_8 is hydrogen, hydrocarbyl, or substituted hydrocarbyl;
- 20 X_9 is an amino protecting group;
 X_{10} is hydrocarbyl, substituted hydrocarbyl, or heteroaryl; and
 Z_2 and Z_4 are independently hydrocarbyl, substituted hydrocarbyl, or heteroaryl.
- 25

13. The process of claim 9 wherein the Lewis acid is selected from the group consisting of the halides or triflates of the Group IB, IIB, IIIB, IVB, VB, VIIB, VIII, IIIA, IVA, lanthanide and actinide elements.

14. The process of claim 13 wherein the Lewis acid is selected from the group consisting of zinc chloride, stannic chloride, cerium trichloride, cuprous chloride, lanthanum trichloride, dysprosium trichloride and ytterbium trichloride.
- 5

15. The process of claim 1 wherein the C(10) acylated taxane comprises a C(7) hydroxy group and the process additionally comprises treating the C(10) acylated taxane with a silylating agent to silylate the
5 C(7) hydroxy group.

16. The process of claim 15 wherein the C(10) acylated taxane is baccatin III.

17. The process of claim 1 wherein the C(10) acylated taxane comprises a C(7) hydroxy group and the process additionally comprises treating the C(10) acylated taxane with an acylating agent to acylate the
5 C(7) hydroxy group.


18. The process of claim 17 wherein the C(10) acylated taxane is baccatin III.

19. The process of claim 1 wherein the C(10) acylated taxane comprises a C(13) hydroxy, metallic oxide, or ammonium oxide substituent and the process additionally comprises the step of esterifying the C(10)
5 acylated taxane by treating the C(10) acylated taxane with a side chain precursor selected from the group consisting of β -lactams, oxazolines, oxazolidine carboxylic acids, oxazolidine carboxylic acid anhydrides, and isoserine derivatives.

20. A process for the silylation of a C(10) hydroxy group of a taxane, the process comprising treating the taxane with a silylamide or a bissilylamide to form a C(10) silylated taxane.

21. The process of claim 20 wherein the silylamide or bissilylamide corresponds to structures 6 or 7, respectively:

77



wherein R_D , R_E , R_F , R_G , and R_H are independently hydrocarbyl, substituted hydrocarbyl, or heteroaryl.

22. The process of claim 21 wherein the taxane is treated with the silylamide or bissilylamide in the presence of an alkali metal base catalyst.

23. The process of claim 21 wherein the taxane is treated with the silylamide or bissilylamide in the presence of a catalyst selected from the group consisting of lithium amide catalysts.


24. The process of claim 20 wherein the silylamide or bissilylamide is selected from the group consisting of tri(hydrocarbyl)silyl-trifluoromethylacetamides and bis tri(hydrocarbyl)-silyltrifluoromethylacetamides, with the hydrocarbyl moiety being substituted or unsubstituted alkyl or aryl.

25. The process of claim 20 wherein the C(10) silylated taxane comprises a C(7) hydroxy group and the process additionally comprises treating the C(10) silylated taxane with an acylating agent to acylate the C(7) hydroxy group.

26. The process of claim 20 wherein the taxane treated with the silylamide or bissilylamide is 10-deacetyl baccatin III and the process additionally comprises treating the C(10) silylated taxane with an acylating agent to acylate the C(7) hydroxy group.


78

27. The process of claim 20 wherein the taxane has the structure:



wherein

- 5 R_1 is hydrogen, hydroxy, protected hydroxy, or together with R_{14} or R_2 forms a carbonate;
 R_2 is keto, $-OT_2$, acyloxy or together with R_1 forms a carbonate;
 R_4 is $-OT_4$ or acyloxy;
 10 R_7 is hydrogen, halogen, $-OT_7$, or acyloxy;
 R_9 is hydrogen, keto, $-OT_9$, or acyloxy;
 R_{10} is hydroxy;
 R_{13} is hydroxy, protected hydroxy, keto, or



- 15 R_{14} is hydrogen, $-OT_{14}$, acyloxy, or together with R_1 forms a carbonate;
 T_2 , T_4 , T_7 , T_9 , and T_{14} are independently hydrogen or hydroxy protecting group;
 X_1 is $-OX_6$, $-SX_7$, or $-NX_8X_9$;
 20 X_2 is hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl;
 X_3 and X_4 are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl;
 X_5 is $-X_{10}$, $-OX_{10}$, $-SX_{10}$, $-NX_8X_{10}$,
 25 or $-SO_2X_{11}$;

X₆ is hydrocarbyl, substituted hydrocarbyl, heteroaryl, hydroxy protecting group or a functional group which increases the water solubility of the taxane derivative;

30 X₇ is hydrocarbyl, substituted hydrocarbyl,
heteroaryl, or sulfhydryl protecting group;

X₈ is hydrogen, hydrocarbyl, or substituted hydrocarbyl;

X₉ is an amino protecting group;

35 X₁₀ is hydrocarbyl, substituted hydrocarbyl, or
heteroaryl;

X₁₁ is hydrocarbyl, substituted hydrocarbyl, heteroaryl, -OX₁₀, or -NX₈X₁₄; and

X₁₄ is hydrogen, hydrocarbyl, substituted
40 hydrocarbyl, or heteroaryl.

28. The process of claim 27 wherein


R₁ is hydroxy or together with R₁₄ or R₂ forms a carbonate;

R₂ is -OCOZ₂, -OCOOZ₂, or together with R₁ forms a
5 carbonate;

R_4 is $-\text{OCOZ}_4$;

R_9 is hydrogen or keto;

R₁₃ is hydroxy, protected hydroxy, or



10 R₁₄ is hydrogen, hydroxy, protected hydroxy, or
together with R₁ forms a carbonate;

$$X_1 \text{ is } -OX_6 \text{ or } -NX_8X_9;$$

X₂ is hydrogen, hydrocarbyl, or substituted hydrocarbyl;


80

- 15 X_3 and X_4 are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl;
 X_5 is $-X_{10}$, $-OX_{10}$, or $-NX_8X_{10}$;
 X_6 is a hydroxy protecting group;
 X_8 is hydrogen, hydrocarbyl, or substituted
20 hydrocarbyl;
 X_9 is an amino protecting group;
 X_{10} is hydrocarbyl, substituted hydrocarbyl, or heteroaryl; and
 Z_2 and Z_4 are independently hydrocarbyl, substituted
25 hydrocarbyl, or heteroaryl.

29. A process for converting the C(7) hydroxy group of a 10-acyloxy-7-hydroxytaxane to an acetal or ketal, the process comprising treating the 10-acyloxy-7-hydroxytaxane with a ketalizing agent in the presence of
5 an acid catalyst to form a C(10) ketalized taxane.

30. The process of claim 29 wherein the C(10) substituent of the 10-acyloxy-7-hydroxytaxane is acetoxy.

31. The process of claim 29 wherein the taxane has the structure:




wherein

- 5 R_1 is hydrogen, hydroxy, protected hydroxy, or together with R_{14} or R_2 forms a carbonate;
 R_2 is keto, $-OT_2$, acyloxy, or together with R_1 forms a carbonate;
 R_4 is $-OT_4$, or acyloxy;

81

- 10 R_7 is hydroxy;
 R_9 is hydrogen, keto, $-OT_9$, or acyloxy;
 R_{10} is acyloxy;
 R_{13} is hydroxy, protected hydroxy, keto, or



- 15 R_{14} is hydrogen, $-OT_4$, acyloxy, or together with R_1 forms a carbonate;
 T_2 , T_4 , T_9 , and T_{14} are independently hydrogen or hydroxy protecting group;
 X_1 is $-OX_6$, $-SX_7$, or $-NX_8X_9$;
20 X_2 is hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl;
 X_3 and X_4 are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl;
 X_5 is $-X_{10}$, $-OX_{10}$, $-SX_{10}$, $-NX_8X_{10}$,
25 or $-SO_2X_{11}$;
 X_6 is hydrocarbyl, substituted hydrocarbyl, heteroaryl, hydroxy protecting group or a functional group which increases the water solubility of the taxane derivative;
30 X_7 is hydrocarbyl, substituted hydrocarbyl, heteroaryl, or sulfhydryl protecting group;
 X_8 is hydrogen, hydrocarbyl, or substituted hydrocarbyl;
 X_9 is an amino protecting group;
35 X_{10} is hydrocarbyl, substituted hydrocarbyl, or heteroaryl;
 X_{11} is hydrocarbyl, substituted hydrocarbyl, heteroaryl, $-OX_{10}$, or $-NX_8X_{14}$; and
 X_{14} is hydrogen, hydrocarbyl, substituted
40 hydrocarbyl, or heteroaryl.

32. The process of claim 31 wherein


R_1 is hydroxy or together with R_{14} or R_2 forms a carbonate;

R_2 is $-\text{OCOZ}_2$, $-\text{OCOOZ}_2$, or together with R_1 forms a
5 carbonate;

R_4 is $-\text{OCOZ}_4$;

R_9 is hydrogen or keto;

R_{13} is hydroxy, protected hydroxy, or



10 R_{14} is hydrogen, hydroxy, protected hydroxy, or together with R_1 forms a carbonate;

X_1 is $-\text{OX}_6$ or $-\text{NX}_8\text{X}_9$;

X_2 is hydrogen, hydrocarbyl, or substituted hydrocarbyl;

15 X_3 and X_4 are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl;

X_5 is $-\text{X}_{10}$, $-\text{OX}_{10}$, or $-\text{NX}_8\text{X}_{10}$;

X_6 is a hydroxy protecting group;

20 X_8 is hydrogen, hydrocarbyl, or substituted hydrocarbyl;

X_9 is an amino protecting group;


X_{10} is hydrocarbyl, substituted hydrocarbyl, or heteroaryl; and

25 Z_2 and Z_4 are independently hydrocarbyl, substituted hydrocarbyl, or heteroaryl.

33. The process of claim 29 wherein the acid catalyst is an inorganic acid.


83

34. The process of claim 29 wherein the ketalizing agent has the formula



wherein X^{31} , X^{32} , X^{33} and X^{34} are independently hydrocarbyl, substituted hydrocarbyl or heteroaryl.


35. The process of claim 29 wherein the ketalizing agent has the formula



wherein X^{31} , X^{32} and X^{33} are independently hydrocarbyl, substituted hydrocarbyl or heteroaryl.

36. The process of claim 29 wherein the ketalizing agent is a vinyl ether.

37. A taxane having the structure:



5

wherein

M is a metal or comprises ammonium:


R_1 is hydrogen, hydroxy, protected hydroxy, or together with R_{14} or R_2 forms a carbonate;

10 R_2 is keto, $-\text{OT}_2$, acyloxy, or together with R_1 forms a carbonate;

- R_4 is $-OT_4$ or acyloxy;
 R_7 is $-OSiR_JR_KR_L$;
 R_9 is hydrogen, keto, $-OT_9$, or acyloxy;
 15 R_{10} is hydrogen, keto, $-OT_{10}$, or acyloxy;
 R_{13} is hydroxy, protected hydroxy, keto, or $MO-$;
 R_{14} is hydrogen, $-OT_{14}$, acyloxy, or together with R_1
 forms a carbonate;
 R_J , R_K , R_L are independently hydrocarbyl, substituted
 20 hydrocarbyl, or heteroaryl, provided, however, if each of
 R_J , R_K and R_L are alkyl, at least one of R_J , R_K and R_L
 comprises a carbon skeleton having at least four carbon
 atoms; and
 T_2 , T_4 , T_9 , T_{10} , and T_{14} are independently hydrogen or
 25 hydroxy protecting group.

38. The taxane of claim 37 wherein R_{10} is $-OT_{10}$, T_{10} is $SiR_D R_E R_F$, and R_D , R_E , R_F are independently hydrocarbyl, substituted hydrocarbyl, or heteroaryl.

39. The taxane of claim 37 wherein
 R_1 is hydroxy or together with R_{14} or R_2 forms a
 carbonate;
 R_2 is $-OCOZ_2$, $-OCOOZ_2$, or together with R_1 forms a
 5 carbonate;
 R_4 is $-OCOZ_4$;
 R_9 is hydrogen or keto;
 R_{13} is hydroxy, protected hydroxy, or



- 10 R_{14} is hydrogen, hydroxy, protected hydroxy, or
 together with R_1 forms a carbonate;
 X_1 is $-OX_6$ or $-NX_8X_9$;

85

X_2 is hydrogen, hydrocarbyl, or substituted hydrocarbyl;

15 X_3 and X_4 are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl;

X_5 is $-X_{10}$, $-OX_{10}$, or $-NX_8X_{10}$;

X_6 is a hydroxy protecting group;


20 X_8 is hydrogen, hydrocarbyl, or substituted hydrocarbyl;

X_9 is an amino protecting group;

X_{10} is hydrocarbyl, substituted hydrocarbyl, or heteroaryl; and

25 Z_2 and Z_4 are independently hydrocarbyl, substituted hydrocarbyl, or heteroaryl.

40. A taxane having the structure:



wherein

M is a metal or comprises ammonium:

5 R_1 is hydrogen, hydroxy, protected hydroxy, or together with R_{14} or R_2 forms a carbonate;

R_2 is keto, $-OT_2$, acyloxy, or together with R_1 forms a carbonate;

R_4 is $-OT_4$ or acyloxy;

10 R_7 is $-OT_7$ or acyloxy;

R_9 is hydrogen, keto, $-OT_9$, or acyloxy;


R_{10} is acyloxy, provided that the acyl moiety is other than acetoxy;

R_{13} is hydroxy, protected hydroxy, keto, or $MO-$;

86

- 15 R_{14} is hydrogen, $-OT_{14}$, acyloxy, or together with R_1 forms a carbonate; and
 T_2 , T_4 , T_7 , T_9 , and T_{14} are independently hydrogen or hydroxy protecting group.

41. A taxane having the structure:



wherein


M is a metal or comprises ammonium:

- 5 R_1 is hydrogen, hydroxy, protected hydroxy, or together with R_{14} or R_2 forms a carbonate;
 R_2 is keto, $-OT_2$, acyloxy, or together with R_1 forms a carbonate;
 R_4 is $-OT_4$ or acyloxy;
 10 R_7 is hydrogen, halogen, $-OT_7$, or acyloxy;
 R_9 is hydrogen, keto, $-OT_9$, or acyloxy;
 R_{10} is $-OSiR_D R_E R_F$;
 R_{13} is hydroxy, protected hydroxy, keto, or $MO-$;
 R_{14} is hydrogen, $-OT_{14}$ or together with R_1 forms a
 15 carbonate;
 R_D , R_E , R_F are independently hydrocarbyl, substituted hydrocarbyl, or heteroaryl; and
 T_2 , T_4 , T_7 , T_9 and T_{14} are independently hydrogen or hydroxy protecting group.

42. The taxane of claim 40 wherein R_7 is acyloxy.

87

43. A taxane having the structure:




wherein

M is a metal or comprises ammonium:

- 5 R_1 is hydrogen, hydroxy, protected hydroxy, or together with R_{14} or R_2 forms a carbonate;
 R_2 is keto, $-OT_2$, acyloxy, or together with R_1 forms a carbonate;
 R_4 is $-OT_4$ or acyloxy;
 10 R_7 is acyloxy;
 R_9 is hydrogen, keto, $-OT_9$, or acyloxy;
 R_{10} is hydrogen, keto, $-OT_{10}$, or acyloxy;
 R_{13} is hydroxy, protected hydroxy, keto, or $MO-$;
 R_{14} is hydrogen, $-OT_{14}$, acyloxy or together with R_1
 15 forms a carbonate; and
 T_2 , T_4 , T_9 , T_{10} and T_{14} are independently hydrogen or hydroxy protecting group.

44. A taxane having the structure:



wherein

M is a metal or comprises ammonium:

- R_1 is hydrogen, hydroxy, protected hydroxy, or together with R_{14} or R_2 forms a carbonate;

88

R_2 is keto, $-OT_2$, acyloxy, or together with R_1 forms a carbonate;

R_4 is $-OT_4$ or acyloxy;

R_7 is $-OT_7$;

R_9 is hydrogen, keto, $-OT_9$, or acyloxy;

R_{10} is hydrogen, keto, $-OT_{10}$, or acyloxy;

R_{13} is hydroxy, protected hydroxy, keto, or $MO-$;

R_{14} is hydrogen, $-OT_{14}$, acyloxy, or together with R_1 forms a carbonate;

T_7 is a ketal or an acetal radical;

T_2 , T_4 , T_9 , T_{10} , and T_{14} are independently hydrogen or hydroxy protecting group.

45. The taxane of claim 44 wherein


R_1 is hydroxy or together with R_{14} or R_2 forms a carbonate;

R_2 is $-OCOZ_2$, $-OCOOZ_2$, or together with R_1 forms a carbonate;

R_4 is $-OCOZ_4$;

R_9 is hydrogen or keto;

R_{13} is hydroxy, protected hydroxy, or



R_{14} is hydrogen, hydroxy, protected hydroxy, or together with R_1 forms a carbonate;

X_1 is $-OX_6$ or $-NX_8X_9$;

X_2 is hydrogen, hydrocarbyl, or substituted hydrocarbyl;

X_3 and X_4 are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl;

X_5 is $-X_{10}$, $-OX_{10}$, or $-NX_8X_{10}$;

X_6 is a hydroxy protecting group;

20 X_8 is hydrogen, hydrocarbyl, or substituted hydrocarbyl;

X_9 is an amino protecting group;

X_{10} is hydrocarbyl, substituted hydrocarbyl, or heteroaryl; and

25 Z_2 and Z_4 are independently hydrocarbyl, substituted hydrocarbyl, or heteroaryl.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/17016

A. CLASSIFICATION OF SUBJECT MATTER		
IPC(6) :C07D 305/14 US CL :549/510, 511 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols) U.S. : 549/510, 511		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5,405,972 A (HOLTON et al.) 11 April 1995, compounds in lines 1-52, col. 25.	37-45
Y	US 5,616,739 A (MAS et al) 01 April 1997, see the entire document.	1-6,15-19, 37-45
Y	DENIS J.-N. et al. A Highly Efficient, Practical Approach to Natural Taxol. J. Am. Chem. Soc. 1988, Vol. 110, pages 5917-5919, see entire document.	1-6,15-19 37-45
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents:	*T*	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		
Date of the actual completion of the international search 11 NOVEMBER 1998	Date of mailing of the international search report 21 DEC 1998	
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer BA K. TRINH <i>for</i> Telephone No. (703) 308-4545	

Form PCT/ISA/210 (second sheet)(July 1992)*